



Pavel Cejnar

**A Condensed Course
of Quantum Mechanics**

KAROLINUM

SECOND EDITION

A Condensed Course of Quantum Mechanics (2nd ed.)

Pavel Cejnar

KAROLINUM PRESS

Karolinum Press is a publishing department of Charles University

Ovocný trh 560/5, 116 36 Prague 1, Czech Republic

www.karolinum.cz, redakcenk@ruk.cuni.cz

© Pavel Cejnar, 2025

Set in the Czech Republic by Karolinum Press

Layout by Jan Šerých

Second edition

A catalogue record for this book is available from the National Library of the Czech Republic.

The original manuscript was reviewed by Jiří Hořejší (Charles University in Prague) and Jean-Paul Blaizot (Paris-Saclay University).

ISBN 978-80-246-6128-5 (pdf)

ISBN 978-80-246-6127-8



Charles University
Karolinum Press

www.karolinum.cz
ebooks@karolinum.cz

Contents

Preface and comments to 1st & 2nd editions. Literature	1
Rough guide to notation	4
Distant outline of quantum physics	6
Introduction	9
Quantum level. Double-slit experiment	9
Wavefunction. Superposition & interference	11
Quantum measurement	13
1a. Space of quantum states	15
Hilbert space & rigged Hilbert space	15
Dirac notation. Linear operators & projectors	18
Sums & products of spaces. Composite systems & entangled states	20
1b. Examples of quantum Hilbert spaces	22
Single structureless particle with spin 0 or $\frac{1}{2}$	22
Ensembles of distinguishable particles	25
Indistinguishable particles. Bosons & fermions. Fock space	26
Artificial systems	30
2a. Representation of quantum observables	32
Operators associated with observables	32
Eigenvalues & eigenvectors in finite & infinite dimension	34
Probability of measurement outcomes	41
2b. Examples of quantum observables	42
Spin- $\frac{1}{2}$ operators	42
Coordinate & momentum	43
Hamiltonian of free particle & particle in potential	44
Orbital angular momentum. Isotropic Hamiltonians	48
Hamiltonian of a particle in electromagnetic field	53
Hamiltonians of simple coupled systems	55
3a. Compatible and incompatible observables	57
Compatible observables. Complete set	57
Incompatible observables. Uncertainty relation	59
Analogy with Poisson brackets	60

Equivalent representations	61
3b. Examples of observable sets	63
Coordinate & momentum. Ladder operators	63
Coordinate & momentum representations	66
Angular momentum operators	68
Addition of two or more angular momenta	71
Complete sets of commuting operators for structureless particle	76
4a. Representation of physical transformations	77
Properties of unitary operators	78
Canonical & symmetry transformations	79
Fundamentals of group theory	82
4b. Examples of symmetry transformations	86
Space translation	86
Space rotation	88
Irreducible representations of the rotation group	92
Spherical tensor operators	94
Space inversion	97
Time translation & reversal. Galilean transformations	98
Symmetry and degeneracy	101
5a. Unitary time evolution	103
Evolution operator & Schrödinger equation	103
Single-particle probability current	104
Conservation laws & symmetries	107
Energy \times time uncertainty	108
Hamiltonians depending on time. Dyson series	112
Schrödinger, Heisenberg & Dirac description	113
Green operator. Single-particle propagator	115
5b. Examples of unitary evolution	118
Two-level system	118
Free particle	119
Coherent states in harmonic oscillator. Ehrenfest theorem	121
Spin in rotating magnetic field	124
6a. Quantum statistical ensembles	126
Density operator. Generalized quantum states	126
Entropy. Canonical ensemble	129

Wigner quasiprobability distribution	131
Density operator for open systems	132
Evolution of density operator: closed & open systems	135
6b. Examples of statistical ensembles	138
Harmonic oscillator at nonzero temperature	138
Coherent superposition vs. statistical mixture	139
Density operator and decoherence for a two-state system	140
7a. Quantum measurement	143
State vector reduction. Interpretation problems	143
A model of measurement. Repeated measurements	146
Measurements on entangled states. EPR situation	149
Quantum nonlocality & Bell inequalities	153
7b. Examples of quantum measurements	156
Destructive & nondestructive measurements	156
Production & measurement of entangled states	158
8. Links between Quantum and Classical	160
Classical limit of quantum theory	160
Feynman integral. Aharonov-Bohm effect. Level density	164
Semiclassical approximation	169
9. Quantum information	174
Quantum information channel	175
Quantum computation	176
10. Stationary approximation methods	181
Variational method	181
Stationary perturbation method	184
Application in atomic physics	189
Application to level dynamics	193
11. Nonstationary approximation methods	197
Nonstationary perturbation method	197
Application to stimulated electromagnetic transitions	203
Driven systems	207
12. Scattering: Iterative approaches	210
Elementary description of elastic scattering	211

General formalism: Lippmann-Schwinger equation	215
Application to elastic scattering	219
13. Scattering: Partial waves	222
Elastic scattering via partial waves	223
Inclusion of inelastic scattering	229
Low-energy & resonance scattering	232
14. Bosonic & Fermionic systems	234
Hilbert space of bosons & fermions	234
Bosonic & fermionic creation/annihilation operators	235
Operators in bosonic & fermionic N -particle spaces	241
Quantization of electromagnetic field	247
15. Many-body techniques	249
Fermionic mean field & Hartree-Fock method	249
Bosonic condensates & Hartree-Bose method	253
Pairing & BCS method	254
Quantum gases	259
Concluding words	266
Index	267

Preface to the first edition

This book was conceived as a collection of notes to my two-semester lecture on quantum mechanics for third-year students of physics at the Faculty of Mathematics and Physics of the Charles University in Prague. It was created in 2011-12.

At first, I just wanted to write down the most important facts, formulas and derivations in a compact form. The information flew in a succinct, “staccato” style, organized in larger and smaller bits (the ■ and ► items), rarely interrupted by wordy explanations. I enjoyed the thick, homogeneous mathematical form of the notes. Calculations, calculations, calculations... I thought of a horrified historian or sociologist who finds no oasis of words. This is how we, tough guys, speak!

However, I discovered that the dense form of the notes was hardly digestible even for tough guys. I had to add some words. To create a “storyteller” who wraps the bare formulas into some minimal amount of phrases. His voice, though still rather laconic, may help to provide the proper motivation and clarify the relevant context. I also formed a system of specific “environments” to facilitate the navigation. In particular: Among crowds of calculations there appears a hierarchy of highlighted formulas:*

important

essential 1

essential 2

crucial

Assumptions or foundational concepts, irreducible to other statements/concepts, appear in boxes:† **Answer to ultimate question of life, universe & everything = 42**

Here and there come some historical notes:‡ **◀ 2013: *Condensed Course* issued**
Handmade schemes (drawn on a whiteboard) illustrate some basic notions.

In this way, the notes have turned into a more serious thing. They almost became a *textbook*! The one distinguished from many others by expanded mathematical derivations (they are mostly given really step by step) and reduced verbal stuffing (just necessary comments in between calculations). This makes the book particularly well suited for conservation purposes—acquired knowledge needs to be stored in a *condensed*, dense enough form, having a compact, nearly tabular structure.

However, as follows from what has been said, this book *cannot* be considered a standard textbook. It may hardly be read with ease and fluency of some more epic treatises. One rather needs to proceed cautiously as a detective, who has to precisely fix all objects on the stage (all symbols, relations etc.) before making any small step forward. This book can be used as a teaching tool, but preferably together with an

*Such formulas are highly recommended to memorize! Although all students of physics & mathematics seem to share a deep contempt for any kind of memorization, I have to stress that all results cannot be rederived in reasonable time limits. There is no escape from saving the key formulas to the memory and using them as quickly reachable starting points for further calculations.

†However, these assumptions do not constitute a closed system of axioms in the strict mathematical sense.

‡I believe that knowledge of history is an important part of understanding. The concepts do not levitate in vacuum but grow from the roots formed by concrete circumstances of their creation. If overlooking these roots, one may misunderstand the concepts.

oral course or a more talkative textbook on quantum mechanics. Below I list some of my favorite candidates for additional guiding texts [1–10].

I have to stress that the notes cover only some parts of *non-relativistic* quantum mechanics. The selection of topics is partly fixed by the settled presentation of the field, and partly results from my personal orientation. The strategy is to introduce the complete general formalism along with its exemplary applications to simple systems (this takes approx. one semester) and then (in the second semester) to proceed to some more specialized problems. Relativistic quantum mechanics is totally absent here; it is postponed as a prelude for the quantum field theory course.

Quantum mechanics is a complex subject. It obligates one to have the skills of a mathematician as well as the thinking of a philosopher. Indeed, the mathematical basis of quantum physics is rather abstract and it is not obvious how to connect it with the observed “reality”. No physical theory but quantum mechanics needs such a sophisticated PR department. We will touch the interpretation issues here, but only very slightly. Those who want to cultivate their opinion (but not to disappear from the intelligible world) are forwarded to the classic [11]. The life saving trick in this *terra incognita* is to tune mind to the joy of thinking rather than to the demand of final answers. The concluding part of the theory may still be missing.

Before we start I should not forget to thank all the brave testers—the first men, mostly students, who have been subject to the influence of this book at its various stages of preparation. They were clever enough to discover a lot of mistakes. Be sure that the remaining mistakes are due to their generous decision to leave some fish for the successors.

In Prague, January 2013

Comments on the second edition

Welcome to the new edition of the *Condensed course*.

While using the first edition for more than a decade of my teaching, I found many items that needed to be fixed, many explanations that should be improved, and many topics that would be worth adding. I have tried to make these important changes in this new edition. In particular, I have made most of the explanations a bit more wordy, I have added several new themes, I have drawn many new figures, I have partly rearranged the content and created a detailed index, and I have corrected numerous misprints.

I hope that the new edition will be much more user-friendly and also more complete than the first one. Though the telegraphic style is deliberately preserved to keep all explications condensed, the book is more viable for all readers, including those with limited initial knowledge. Extensions and new topics make the book more robust, providing necessary initial knowledge for most of the main presently active directions of nonrelativistic quantum theory. I believe that the *Condensed*

course in the present form offers a balanced concise introduction to the traditional topics, related to the general formalism and natural quantum systems, as well as to modern topics, focused on artificial quantum systems and quantum information.

And the last but not least: I pay off my big debt from the first edition by adding the following “essential historical remark”:

◀ Essential historical remark ☺

1902: Jára Cimrman anticipates quantum uncertainty by studying his rat trap bait-box mechanism & answering naughty teenager’s questions of E. Schrödinger

In Prague, August 2025

Recommended textbooks:

- [1] J.J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, 1985, 1994)
- [2] J.J. Sakurai, J.J. Napolitano, *Modern Quantum Mechanics* (Addison-Wesley, 2011)
(a modified edition of [1])
- [3] G. Auletta, M. Fortunato, G. Parisi, *Quantum Mechanics* (Cambridge University Press, 2009)
- [4] L.E. Ballentine, *Quantum Mechanics. A Modern Development* (World Scientific, Singapore, 1998)
- [5] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, 1995)
- [6] A. Bohm, *Quantum Mechanics: Foundations and Applications* (Springer, 1979, 1993)
- [7] W. Greiner, *Quantum Mechanics: An Introduction* (Springer, 1989)
W. Greiner, *Quantum Mechanics: Special Chapters* (Springer, 1998)
W. Greiner, B Müller, *Quantum Mechanics: Symmetries* (Springer, 1989)
- [8] E. Merzbacher, *Quantum Mechanics* (Wiley, 1998)
- [9] V. Zelevinsky, *Quantum Physics*, Volume 1 & 2 (Wiley-VCH, 2011)
- [10] A. Messiah, *Quantum Mechanics* (Dover, 1999)
(living classic, first published in 1958)

Further reading:

- [11] J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, 1987)
(a collection on brilliant essays on the interpretation of quantum theory)
- [12] R. Omnès, *The Interpretation of Quantum Mechanics* (Princeton University Press, 1994)
(a more systematic treatment of the interpretation questions)
- [13] T. Lancaster, S.J. Blundell, *Quantum Field Theory for a Gifted Amateur* (Oxford Univ. Press, 2014)
(a readable introduction to the world behind nonrelativistic QM)
- [14] D. Griffiths, *Introduction to Elementary Particles* (Wiley-VCH, 2008)
(an accessible overview of the standard model of fundamental particles and interactions)
- [15] C. Gardiner, P. Zoller *The quantum world of ultracold atoms and light*
Book I: *Foundations of quantum optics* (Imperial College Press, 2014)
Book II: *The physics of quantum-optical devices* (Imperial College Press, 2015)
Book III: *Ultra-cold atoms* (World Scientific, 2017)
(an introduction to controllable quantum systems)
- [16] A. Pais, *Inward Bound of Matter and Forces in the Physical World* (Clarendon Press, 1986)
(an exciting treatise on the history of the physics of microworld)

Rough guide to notation (no notation is perfect!)

$\alpha \psi\rangle + \beta \psi'\rangle$ $\{ \phi_i\rangle\}_{i=1}^{d_{\mathcal{H}}}, d_{\mathcal{H}}$ $ \psi\rangle, \langle\psi' , \langle\psi' \psi\rangle$ $\ \psi\ = \sqrt{\langle\psi \psi\rangle} = 1/\mathcal{N}$ $\underline{\mathcal{H}}, \mathcal{H}, \tilde{\mathcal{H}}$ $\ell^2, \mathcal{L}^2(\mathbb{R}^3), \mathbb{C}^d$ $\text{Span}\{ \psi_1\rangle \dots \psi_n\rangle\}$ $\mathcal{H}^{(N)}, \mathcal{H}_{\pm}^{(N)}$ $\mathcal{H}_1 \otimes \mathcal{H}_2, \bigotimes_{i=1}^n \mathcal{H}_i, \mathcal{H}_1 \oplus \mathcal{H}_2, \bigoplus_{i=1}^n \mathcal{H}_i$ $ \psi\rangle_1 \psi'\rangle_2, \Phi_{ij}\rangle \equiv \phi_{1i}\rangle_1 \phi_{2j}\rangle_2$ $\psi(\vec{x}) \equiv \langle \vec{x} \psi \rangle, \tilde{\psi}(\vec{p}) \equiv \langle \vec{p} \psi \rangle$ $\psi(\vec{x}, m_s) \equiv \Psi(\vec{x})$ $\Psi(\xi_1 \dots \xi_N)$ $ a\rangle, a^{(k)}\rangle, a_i\rangle, a_i^{(k)}\rangle$ $ E_i\rangle, E_i^{(k)}\rangle, E\rangle$ $ \uparrow\rangle, \downarrow\rangle$ $ lm\rangle, sm_s\rangle, jm_j\rangle$ $R_{nl}(r) = u_{nl}(r)/r, R_{kl}(r)$ $C_{j_1 m_1 j_2 m_2}^{j m} \equiv \langle j_1 j_2 j m j_1 m_1 j_2 m_2 \rangle$ $ \psi_{ni}\rangle, \psi_i^{(n)}(\lambda)\rangle$ $ 0\rangle, n_1, n_2, \dots\rangle$ $ \Psi_{\text{HF}}\rangle, \Psi_{\text{HB}}\rangle, \Psi_{\text{BCS}}\rangle$	<p>Hilbert spaces, vectors & wavefunctions, scalar products</p> <p>superposition \equiv linear combination of state vectors ($\alpha, \beta \in \mathbb{C}$)</p> <p>general set of basis vectors in Hilbert space \mathcal{H}, dimension of \mathcal{H}</p> <p>ket & bra forms of state vectors, scalar product</p> <p>norm of vector = 1/normalization coefficient</p> <p>Gelfand's hierarchy of spaces (rigged Hilbert space)</p> <p>specific separable or finite Hilbert spaces</p> <p>linear space spanned by the given vectors</p> <p>N-particle Hilbert space, its exchange symmetric/antisym.subspaces</p> <p>direct product & sum of Hilbert spaces</p> <p>general factorized state vector, factorized basis in $\mathcal{H}_1 \otimes \mathcal{H}_2$</p> <p>wavefunction of spinless particle in coordinate & momentum repres.</p> <p>single-particle wavefunction in single/multicomponent forms</p> <p>N-particle wavefunction with $\xi_1 \equiv (\vec{x}_i, m_i)$</p> <p>eigenvector of operator \hat{A} with eigenvalue a or a_i (degeneracy index k)</p> <p>eigenvectors of Hamiltonian (discrete or continuous energy)</p> <p>up & down projection states of spin $s = \frac{1}{2}$</p> <p>states with orbital, spin & total ang. momentum l, s & j, projection m.</p> <p>radial wavefunction ($n \equiv$ princ.q.num., $l \equiv$ orb.ang.mom., $k \equiv$ wave vec.)</p> <p>Clebsch-Gordan coefficient for the coupling of 2 angular momenta</p> <p>nth-order perturbation correction & approx. of ith energy eigenstate</p> <p>vacuum state, basis states of $\mathcal{H}_{\pm}^{(N)}$ in occupation-number repres.</p> <p>Hartree-Fock/Bose & BCS approx. of many-body ground state</p>
$\hat{O}, \hat{O}^\dagger, \hat{O}^{-1}$ $O_{ij} = \langle \phi_i \hat{O} \phi_j \rangle$ $\ \hat{O}\ , \text{Def}(\hat{O})$ $\hat{A}, \hat{U}, \hat{I}, \hat{I}_{\mathcal{H}}$ $\hat{A}_{\text{S}}, \hat{A}_{\text{H}}(t), \hat{A}_{\text{D}}(t)$ \hat{A} $\hat{A}_1 \otimes \hat{A}_2$ $\mathcal{S}(\hat{A}), \mathcal{D}(\hat{A}), \mathcal{C}(\hat{A})$ $\hat{P}_0, \hat{P}_{\pm}^{(N)}$ $\hat{P}_a, \hat{\Pi}_a, \hat{\Pi}_{(a_1, a_2)}$ $\vec{\nabla}, \Delta$ $\hat{x}, \hat{p}, \hat{\pi}$ $\hat{H}, \hat{K}, \hat{V}, \hat{H}'$ $\hat{L}, \hat{S}, \hat{J}$ $\hat{J}_0 \equiv \hat{J}_z, \hat{J}_{\pm} \equiv \hat{J}_x \pm i\hat{J}_y$ $\hat{\sigma} \equiv (\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z)$ $\hat{D}, \hat{\mu}$ $\hat{b}_k, \hat{b}_k^\dagger; \hat{a}_k, \hat{a}_k^\dagger; \hat{c}_k, \hat{c}_k^\dagger$ \hat{N}, \hat{N}_k $\hat{O}^{(n)}$ $\hat{T}_{\vec{a}}, \hat{T}_{\Delta o}$ $\hat{R}_{\vec{n}\phi} \equiv \hat{R}_{\mathbf{R}}, \mathbf{R}(\alpha\beta\gamma)$ $\hat{P}, \hat{\mathcal{T}}$ $\hat{G}_i, \hat{C}_{\mathcal{G}}$	<p>Operators: observables, transformations & evolution</p> <p>linear operator, its Hermitian conjugate & inverse</p> <p>matrix element of operator \hat{O}</p> <p>norm & definition domain of operator</p> <p>general Hermitian & unitary operator, identity operator (in space \mathcal{H})</p> <p>Schrödinger, Heisenberg, Dirac representations of observable</p> <p>operator expressing time derivative of observable</p> <p>tensor product of operators acting in $\mathcal{H}_1 \otimes \mathcal{H}_2$</p> <p>full spectrum of observable \hat{A}, its discrete & continuous parts</p> <p>projector to a general subspace $\mathcal{H}_0 \subset \mathcal{H}$, projector to $\mathcal{H}_{\pm}^{(N)}$</p> <p>projectors to discrete & continuous eigenvalue subspaces</p> <p>gradient & Laplace operator (if not an interval or gap)</p> <p>coordinate operator, canonical & mechanical momentum operator</p> <p>Hamiltonian, its kinetic & potential terms, Hamiltonian perturbation</p> <p>orbital, spin & total angular momentum operators</p> <p>spherical components of \hat{J}, shift operators for $jm\rangle$ eigenstates</p> <p>the triplet of Pauli matrices</p> <p>operators of electric & magnetic dipole moments</p> <p>annih. & creation operator of boson, fermion or gen.particle in state $\phi_k\rangle$</p> <p>total number of particles & number of particles in basis state $\phi_k\rangle$</p> <p>n-body operator</p> <p>space translation or general eigenstate shift operator $o\rangle \rightarrow o + \Delta o\rangle$</p> <p>rotation operator in \mathcal{H} (axis, angle) & rot.matrix in 3D (Euler angles)</p> <p>space inversion operator (parity) & time reversal operator</p> <p>generator & Casimir operator of a group \mathcal{G}</p>

$\hat{U}(t), \hat{U}(t_1, t_0)$ $\hat{G}^\pm(t, t_0), G^+(\vec{x}t \vec{x}_0t_0)$ $\hat{G}^\pm(E), \hat{T}^\pm(E)$ $\langle \Phi_{E'n'} \hat{S} \Phi_{En} \rangle, \langle \phi_{\vec{k}'} \hat{S} \phi_{\vec{k}} \rangle$ \mathfrak{T} $[\hat{A}^{\lambda_1} \times \hat{B}^{\lambda_1}]_\mu^\lambda$ $[\hat{A}, \hat{B}], \{\hat{A}, \hat{B}\}$ $\{A, B\}$ $\text{Tr } \hat{O}, \text{Tr}_1 \hat{O}, \text{Det } \hat{O}$	evolution operator for times $t_0 \xrightarrow{t} t_1$ retarded & advanced Green operators, propagator energy image of Green operators, T -operator in scattering theory S-matrix elements time ordering of operator product tensor coupling of spherical tensor operators $\hat{A}_{\mu_1}^{\lambda_1}, \hat{B}_{\mu_2}^{\lambda_1}$ commutator & anticommutator of operators Poisson bracket of classical observables trace of operator/matrix, partial trace over \mathcal{H}_1 in $\mathcal{H}_1 \otimes \mathcal{H}_2$, determinant
$\alpha_\psi(\psi'), \alpha_\psi(x)$ $\mathfrak{p}_\psi(\psi'), \mathfrak{p}_\psi(\mathcal{H}_0)$ $\mathfrak{p}_\psi(x)$ $\alpha_0(t), \mathfrak{p}_0(t)$ $\mathfrak{p}_c(a b)$ $\alpha_{ji}(t), \mathfrak{p}_{ji}(t), \mathcal{R}_{ji}(t), \mathcal{R}_X$ $\langle A \rangle_\psi, \langle a \rangle_c$ $\langle\langle A^2 \rangle\rangle_\psi \equiv \Delta_\psi^2 A$ $\rho(\vec{x}, t), \vec{j}(\vec{x}, t)$ $\hat{\rho}(t), \hat{\rho}_1(t) = \text{Tr}_2 \hat{\rho}(t)$ $W_\rho(\vec{x}, \vec{p}, t)$ $\rho(\vec{x}, \vec{p}, t)$ $\varrho(E), \varrho_f(E), \varrho(\xi)$	Statistics, probabilities & densities amplitude to identify $ \psi\rangle$ with $ \psi'\rangle$ or to measure value x of an observable probability to identify $ \psi\rangle$ with $ \psi'\rangle$ or with an arbitrary state from $\mathcal{H}_0 \subset \mathcal{H}$ probability to measure values x of some observables in state $ \psi\rangle$ survival amplitude & probability of $t=0$ initial state at time t conditional probability of a given b (depending on parameter c) $ \phi_i\rangle \xrightarrow{t} \phi_j\rangle$ transition amplitude, probability & rate, rate of event X average value of observable A in $ \psi\rangle$, average of a for a fixed parameter c variance of the distribution $\mathfrak{p}_\psi(a)$ (squared uncertainty of observable A) single-particle probability density & flow at point \vec{x} , time t general density operator, density operator of a subsystem (partial trace) Wigner quasiprobability distribution in phase space for a given $\hat{\rho}$ classical probability distribution in phase space level density, density of final states, particle density at $\xi \equiv (\vec{x}, m_s)$
$\hbar = h/2\pi$ c, e, ϵ_0, α $\lambda_C, \lambda_C, \lambda_B, \lambda_B, a_B$ \vec{k}, ω M, \mathcal{M}, q $E, E_i, E_{ni}, E_i^{(n)}(\lambda)$ ε_k, n_k $V, \vec{A}, \vec{E}, \vec{B}$ $\frac{d\sigma}{d\Omega}, \sigma^{\text{el}}, \sigma^{\text{inel}}, \sigma^{\text{tot}}$ $f_{\vec{k}}(\vec{k}'), f_{n\vec{k}}(\vec{k}'), f_{\vec{k}}^{(n)}(\vec{k}')$ $F_l(k), S_l(k), \delta_l(k), \eta_l(k)$ R, l_{max} S_ρ $Z(\beta), Z(\beta, \mu)$ $S[\vec{x}(t)], S(\vec{x}, t), \mathcal{L}(\vec{x}, \dot{\vec{x}})$	Physical constants & parameters, various physical quantities reduced & unreduced Planck constant speed of light, elementary charge, vacuum permittivity, fine-structure const. reduced & unreduced Compton & de Broglie wavelengths, Bohr radius wavevector, angular frequency particle mass, two-particle reduced mass, particle charge continuous & discrete energy, its n^{th} order perturb. correction & approximation energies & occupation numbers of single-particle states scalar & vector electromagnetic potentials, el. intensity & mag. induction differential cross section, integral elastic, inelastic & total cross sections scattering amplitude, its n^{th} order Born correction & approximation partial wave amplitude, S-matrix, phase shift & inelastic suppression factor range of potential, maximal orbital angular momentum von Neumann entropy of density operator $\hat{\rho}$ (grand)canonical partition function ($\beta \equiv$ inverse temp., $\mu \equiv$ chem. pot.) classical action (functional & function forms), Lagrangian
$j_l, n_l, h_l^\pm(kr)$ $L_l^j(\rho), H_n(\xi)$ $P_{lm}(\cos \vartheta), Y_{lm}(\vartheta, \varphi)$ $D_{m'm}^j(\alpha\beta\gamma) \equiv D_{m'm}^j(\mathbf{R})$ $\delta(x), \delta_\epsilon(x), \Theta(x)$ $\delta_{ij}, \varepsilon_{ijk}$ $(1, 2, 3) \equiv (x, y, z)$ $\vec{n}, \left\{ \begin{array}{l} (\vec{n}_x, \vec{n}_y, \vec{n}_z) \\ (\vec{n}_r, \vec{n}_\vartheta, \vec{n}_\varphi) \end{array} \right\}$ $\{X_i\}_{i=1}^n, \{X_i\}_{i \in \mathcal{D}}, \{X(c)\}_{c \in \mathcal{C}}$ Min, Max, Sup $\{X_i\}_i$ iff, l.h.s., r.h.s	Special functions & miscellaneous mathematical symbols Bessel, Neumann & Hankel functions associated or generalized Laguerre polynomials & Hermite polynomials associated Legendre polynomial, spherical harmonics ($\vartheta, \varphi \equiv$ sph. angles) Wigner matrix/function (Euler angles of rotation matrix) Dirac δ -function, imperfect δ functions, step function Kronecker & Levi-Civita symbols indices of Cartesian components unit vector, $\left\{ \begin{array}{l} \text{Cartesian} \\ \text{spherical} \end{array} \right\}$ orthonormal coordinate vectors discrete/continuous set of objects minimum, maximum, supremum of a set of numbers “if and only if”, the left- / right-hand side (of an equation)

Distant outline of quantum physics

Historical origins: Quantum mechanics was born in the 1900s in analyses of (i) electromagnetic radiation emitted by matter in thermal equilibrium and (ii) specific heats of solids at low absolute temperatures. A few years later, the discovery of the structure of atom implied a more fundamental problem: (iii) the question of stability of matter. A solution of all these problems was found in a modification of the laws of classical (Newtonian) physics by assuming some particular rules of quantization for certain physical quantities like energy. These principles (which invited the word “quantum”) moreover explained an older mystery of discrete spectra of light radiated by single elements. However, it turned out that a much more radical modification of the physics paradigm was needed. The consistent theory of quantum phenomena was build in the piece by piece manner during the 1920s and 1930s. This development explains why quantum theory (in contrast to Einstein’s relativity) carries traces of rather different approaches and ways of thinking. Discussions on the interpretation of quantum theory continue up to the present days.

Probabilistic character: Quantum physics is ultimately indeterministic. It does not generally predict precise outcomes of individual experiments but only probabilities of various alternative results. It is the only theory in which randomness represents a really fundamental concept (its use in the classical context is just a tool to overcome a lack of information). Quantum physics may be considered as a simultaneous description of multiple alternatives of physical reality with no possibility to predict which of the alternatives will be finally actualized for a particular observer.

Linearity: Underlying the dynamics of quantum probabilities, there is a rather simple linear theory which makes use of so-called quantum amplitudes. An amplitude $\mathbf{a} = |\mathbf{a}|e^{i\varphi}$ of a certain physical event is a number inside the unit circle of the \mathbb{C} plane such that the probability of the event is $\mathbf{p} = |\mathbf{a}|^2$. Though the observable output (probability \mathbf{p}) is contained only in $|\mathbf{a}|$, the phase angle φ is irreducible. Manifestation of linearity is twofold: (i) If a given system can be prepared in two particular initial states, denoted as $|\psi_1\rangle$ and $|\psi_2\rangle$ (generalization to more states is obvious), quantum theory requires that it can also be prepared in a state $\alpha_1|\psi_1\rangle + \alpha_2|\psi_2\rangle$, which corresponds to a linear combination (quantum superposition) of the above two states with arbitrary complex coefficients α_1 and α_2 . The meaning of quantum superpositions is highly counterintuitive — e.g., they may represent states in which a particle simultaneously takes several positions. (ii) If the quantum amplitudes of a given measurement outcome for the two initial states are \mathbf{a}_1 and \mathbf{a}_2 , the corresponding amplitude for the above superposition is $\mathbf{a} = \mathcal{N}(\alpha_1\mathbf{a}_1 + \alpha_2\mathbf{a}_2)$, where the normalization coefficient $\mathcal{N} \in \mathbb{R}$ ensures that an integral of $\mathbf{p} = |\mathbf{a}|^2$ over all possible outcomes is equal to 1. Linearity of amplitudes implies nonlinearity of probabilities, which is the key for explanation of various quantum interference effects.

Complementarity: In quantum theory, all conceivable quantities that can be measured on a given system are sorted according to their mutual compatibility. Any observable is compatible only with a subset of the remaining observables and incompatible with the others. Any set of compatible observables can be simultaneously known with certainty, but this knowledge excludes a precise determination of any incompatible observable. Joint probabilities of simultaneous measurement outcomes can be consistently determined only for sets of compatible observables; for sets of incompatible observables they depend on details of the measurement procedure.

Nonseparability: Evolution of a given quantum system S often includes interaction with an external environment and/or other degrees of freedom E. Linearity of quantum theory leads to creation of superpositions of the composite system S + E that have a form $\sum_i \alpha_i |\psi_i\rangle_S |\psi'_i\rangle_E$. Here $|\psi_i\rangle_S$ and $|\psi'_i\rangle_E$ are mutually correlated states of S and E, joint into a separable state $|\psi_i\rangle_S |\psi'_i\rangle_E$ of the S + E system, and α_i are some coefficients. The whole superposition (unlike its individual terms) cannot in general be factorized to a single product of S and E states. Hence in these so-called entangled states the subsystems S and E are not separable. An ensemble of interacting quantum subsystems can become a strongly holistic object in which correlations between distant parts are stronger than allowed in classical physics.

Quantum measurement: The entanglement process takes place also during the act of a general measurement. Unfactorizable superpositions resulting from this process correlate various states of the measuring apparatus (different measurement outputs) with the associated states of the measured system. Identifying the actual reality with only a single output, we select only a part of the superposition. This is often treated as an irreducible influence of quantum measurement (or of an observer, who may be considered as the “selector” of reality) on the measured object.

Links to other branches of physics: Quantum physics is a continuation of classical physics to the world of small objects and/or tiny actions. It is treated in two parts: the nonrelativistic and relativistic quantum theory. Since the combination of relativistic and quantum laws implies new phenomena, the general formalism of quantum theory is first applied to nonrelativistic mechanics, which is sufficient in the description of a large class of objects. The same formalism is subsequently recalled in the context of special relativity, leading to the quantum field theory, which provides so far the deepest description of elementary particles of matter and their mutual interactions. Unification of quantum theory with general relativity (theory of gravity) is not available yet. Quantum theory is a basis for great majority of contemporary “applied” physics, like molecular, atomic, nuclear and subnuclear physics, condensed matter and solid-state physics, optics, astrophysics etc. Recently, some particular applications of quantum laws gave rise to a special branch of physics called “quantum information”.



HEALTH AUTHORITY WARNING
THINKING ABOUT
QUANTUM PHYSICS
CAUSES INSOMNIA

INTRODUCTION

Before sailing out, we encourage the crew to get ready for adventures. Quantum mechanics deals with phenomena, which are rather unusual from the viewpoint of our common macroscopic experience. Description of these phenomena makes us sacrifice some principles which we used to consider self-evident.

■ Quantum level

Quantum theory describes objects on the atomic and subatomic scales, but also larger objects if they are observed with an extremely **high resolution**.

► Planck constant

The domain of applicability of quantum mechanics is determined by constant


$\hbar \doteq 1.05 \cdot 10^{-34} \text{ J}\cdot\text{s} \doteq 0.66 \text{ eV}\cdot\text{fs}$ which defines a **quantum unit of action**

► Phenomena whose actions are on/below the scale of \hbar belong to the quantum jurisdiction. However, even phenomena with larger absolute actions can get to the quantum domain if the difference of actions between distinguishable alternatives reaches the \hbar scale. Consider two trajectories $\mathbf{q}_1(t)$ & $\mathbf{q}_2(t)$ in the configuration space of the system (\mathbf{q} is a multidimensional vector of generalized coordinates depending on time t) which, in the given experimental situation, are on the limit of mutual distinguishability (so these and similar trajectories can still be experimentally distinguished from each other, but the trajectories which are closer than these cannot). The classical action of each trajectory is $S[\mathbf{q}_\bullet(t)]$. The difference $\Delta S = |S[\mathbf{q}_1(t)] - S[\mathbf{q}_2(t)]|$ determines whether the situation can be described in the classical or quantum way:

Classical mechanics Quantum mechanics	} applies if the difference satisfies	$\Delta S \gg \hbar$ $\Delta S \lesssim \hbar$
--	---------------------------------------	---

In particular, if the minimum of the action functional S expressed on the level of resolution $\Delta S \sim \hbar$ extends across several distinguishable trajectories, all these trajectories must be *somehow* taken into account *simultaneously*.

Quantum description is then unavoidable.



◀ Historical remark

1900: Max Planck introduced \hbar along with the quanta of electromagnetic radiation to explain the blackbody radiation law

1905: Albert Einstein confirmed elmag. quanta in the explanation of photoeffect


1913: Niels Bohr introduces a quantum model of atoms (“old quantum mechanics”)

■ Double-slit experiment


According to Richard Feynman and many others, this is the most crucial quantum experiment that allows one to realize how unusual the quantum world is. There exist numerous variations and improvements of this experiment.

► Arrangement

Components: Emitter E which emits particles (in the one by one mode), the plate with open slits A and B, the screen S where positions of arriving particles are detected (dots)



Both particle trajectories $\vec{x}_A(t)$ and $\vec{x}_B(t)$ from the emitter (\vec{x}_E) to the screen (\vec{x}_S) minimize the action functional $S[\vec{x}(t)]$.
Suppose $|S_A - S_B| \lesssim \hbar$



► Regimes and results of measurements

(a) **Interference setup:** position of the particle is measured only at the screen \Rightarrow individual particle hits are randomly scattered within strips that form a wave-like interference pattern

(b) **Which-path setup:** prior the screen measurement, the particle position is measured—either explicitly (with the results observed), or implicitly (results hidden)—immediately after the slits \Rightarrow individual particle hits at the screen cumulate straight behind the slits, no interference behavior is observed

Delayed choice: The choice of setup (a)/(b) is made *after* the particle passed the slits. The outcome is the same as if the decision was made before.

Quantum eraser: The unobserved which-path information from setup (b) is erased before the particle hits the screen. The interference pattern appears.

► Some conceptual implications

Indeterminism: It is not possible to predict the positions of individual particle hits, but only their overall distribution. Quantum physics invites randomness and probabilistic description into the fundamental theory.

Particle-wave duality: Particles show either wave or corpuscular properties, in accord with the specific experimental arrangement. In particular, the existence of the which-path information invariably leads to the corpuscular behavior, while its actual nonexistence implies a wave-like behavior.

Contextuality etc.: The actual result of a physical observation depends on a wider “context” of the process investigated. The observed “reality” emerges only during the act of observation. And many more sentences like these.

◀ Historical remark

1805 (approx.): Thomas Young performed double-slit experiment with light

1927: C. Davisson & L. Germer demonstrate interference of electrons on crystals

1961: first double-slit experiment with massive particles (electrons)

1970’s: double-slit experiments with individual electrons

1990’s-present: progress in realizations of which-path setup & delayed-choice exp.


■ Wavefunction and superposition principle

To explain the outcome of the interference setup of the double-slit experiment (interference pattern formed by individual dots), we will assume that the undisturbed particle inside the interferometer represents a wave-like object comprising a variety of potential particle localizations and that the position measurement on the screen makes one of these potential localizations actual.

► Concept of wavefunction

Quantum physics deals not with one, but with **several alternative versions of reality**—with many potential outcomes of any conceivable measurement performed on a given system. Complete determination of the physical state of the system must somehow include all these alternatives and to quantify their probabilities. If variable x denotes possible outcomes of a complete set of measurements (specifying all degrees of freedom of a given system), the quantum state of the system is determined by a complex wavefunction $\psi(x)$:

Wavefunction value $\boxed{\psi(x) \equiv \text{amplitude}}$
of probability (or density of amplitude
of probability if x is continuous) for
finding the particular alternative x .



Squared modulus $\boxed{|\psi(x)|^2 \equiv \text{probability}}$

(or density of probability for x continuous) for finding the alternative x .

Although the detectable probabilities are given by $|\psi(x)|^2 \in \mathbb{R}$, their amplitudes $\boxed{\psi(x) \in \mathbb{C}}$ play a substantial role in the quantum description of reality!

The wavefunction evolves in time t , so: $\psi(x) \rightarrow \psi(x, t)$

► Wavefunction of a single **structureless particle**: $\boxed{\psi(\vec{x}, t) \equiv \sqrt{\rho(\vec{x}, t)} e^{i\varphi(\vec{x}, t)}}$

where $\vec{x} \equiv$ alternative positions of the particle in the real 3D space

$|\psi(\vec{x}, t)|^2 = \rho(\vec{x}, t) \geq 0$ is the probability density to detect the particle at position \vec{x} . Normalization: $\int \rho(\vec{x}, t) d\vec{x} = 1 \forall t$. Phase $\varphi(\vec{x}, t) \in \mathbb{R}$ has no “classical” interpretation, but plays an important role in interference phenomena

► Superposition of wavefunctions

The outcome of the interference setup depends on the fact that waves can be summed up. Consider two normalizable wavefunctions $\psi_A(\vec{x}, t)$ and $\psi_B(\vec{x}, t)$:

$$\int |\psi_A|^2 d\vec{x} < \infty, \int |\psi_B|^2 d\vec{x} < \infty \Rightarrow \boxed{\int |\alpha\psi_A + \beta\psi_B|^2 d\vec{x} < \infty} \quad \forall \alpha, \beta \in \mathbb{C}$$

\Rightarrow any linear combination of normalizable wavefunctions is a normalizable wavefunction \Rightarrow these functions form a linear vector space $\mathcal{L}^2(\mathbb{R}^3)$

► Interference phenomenon

Probability density for a superposition of waves is not the sum of densities for individual waves. Choose arbitrary $\alpha = |\alpha|e^{i\varphi_\alpha}$ and $\beta = |\beta|e^{i\varphi_\beta}$ such that $\int |\alpha\psi_A + \beta\psi_B|^2 d\vec{x} = 1$ with both ψ_A and ψ_B normalized ($\int |\psi_\bullet|^2 d\vec{x} = 1$)

$$\Rightarrow \boxed{\underbrace{|\alpha\psi_A + \beta\psi_B|^2}_{\rho_{\alpha A + \beta B}} = \underbrace{|\alpha\psi_A|^2}_{|\alpha|^2 \rho_A} + \underbrace{|\beta\psi_B|^2}_{|\beta|^2 \rho_B} + \underbrace{2|\alpha\beta\psi_A\psi_B| \cos(\varphi_A + \varphi_\alpha - \varphi_B - \varphi_\beta)}_{\text{interference terms}}}$$

► Description of the interference setup in the double-slit experiment

Despite generally delocalized nature of wavefunctions we assume an approximate assignment of times: at $t \approx t_0$ the particle passes the double-slit plate and at $t \approx t_1$ it reaches the detection screen. At the plate we have

$$\psi(\vec{x}, t_0) \approx \alpha\delta_A(\vec{x} - \vec{x}_A) + \beta\delta_B(\vec{x} - \vec{x}_B)$$

with $\delta_\bullet(\vec{x} - \vec{x}_\bullet)$ denoting the wavefunction localized at the respective slit ($\delta_\bullet = 0$ away from it) and α, β some coefficients depending on the emitted state and experimental details. If $\psi_\bullet(\vec{x}, \Delta t)$ is the wavefunction developed in time $\Delta t = t_1 - t_0$ from $\delta_\bullet(\vec{x} - \vec{x}_\bullet)$, the wavefunction on the screen reads as:


$$\psi(\vec{x}, t_1) \approx \alpha\psi_A(\vec{x}, \Delta t) + \beta\psi_B(\vec{x}, \Delta t) \quad \Rightarrow \quad \boxed{\rho(\vec{x}) \approx |\alpha\psi_A(\vec{x}, \Delta t) + \beta\psi_B(\vec{x}, \Delta t)|^2}$$

Thus the probability distribution on the screen shows the interference pattern.

► Dirac delta function (mathematical intermezzo)

To deal with arbitrary wavefunctions, it is convenient to introduce a generalized function (more precisely, a so-called distribution) describing a perfectly localized particle. Consider first the 1D case. In a vague sense, the δ -function can be seen as a “limit” of a series of ordinary functions whose support contracts to a single point but the integral remains constant, equal to unity:

$$\delta(x) = \lim_{\epsilon \rightarrow 0} \delta_\epsilon(x) \quad \text{Support}[\delta(x)] \equiv \{x=0\} \quad \text{and} \quad \int_{-\infty}^{+\infty} \delta(x) dx = 1$$




For instance, we can choose the following sequences:

(a) $\delta_\epsilon(x) \equiv \begin{cases} \frac{1}{\epsilon} & \text{for } x \in [-\frac{\epsilon}{2}, +\frac{\epsilon}{2}] \\ 0 & \text{otherwise} \end{cases}$

(b) $\delta_\epsilon(x) = \frac{1}{\pi} \frac{\epsilon}{\epsilon^2 + x^2}$ (Cauchy or Breit-Wigner form)

(c) $\delta_\epsilon(x) = \frac{1}{\sqrt{2\pi\epsilon^2}} e^{-\frac{x^2}{2\epsilon^2}}$ (Gaussian form)

(d) $\delta_\epsilon(x) = \frac{1}{\pi} \frac{\sin(x\epsilon^{-1})}{x} = \frac{1}{2\pi} \int_{-\epsilon^{-1}}^{+\epsilon^{-1}} e^{iqx} dq$ (Fourier transform of unity)



In 3D space:

$$\overbrace{\delta_{\epsilon_1}(x_1 - x'_1) \delta_{\epsilon_2}(x_2 - x'_2) \delta_{\epsilon_3}(x_3 - x'_3)}^{\delta_\epsilon(\vec{x} - \vec{x}')} \xrightarrow{\epsilon \rightarrow 0} \overbrace{\delta(x_1 - x'_1) \delta(x_2 - x'_2) \delta(x_3 - x'_3)}^{\delta(\vec{x} - \vec{x}')}$$

Defining property of δ -function in terms of distribution theory:

$$\int f(\vec{x}) \delta(\vec{x} - \vec{x}') d\vec{x} = f(\vec{x}')$$

► **Delocalized wavefunctions**

Any wavefunction can be expressed as: $\psi(\vec{x}, t) = \int \psi(\vec{x}', t) \delta(\vec{x} - \vec{x}') d\vec{x}'$

General state given by a wavefunction $\psi(\vec{x}, t) \equiv$ **superposition of localized states** $\delta(\vec{x} - \vec{x}')$ with coefficients equal to the respective values $\psi(\vec{x}', t)$

However, note that $\delta(\vec{x} - \vec{x}') \notin \mathcal{L}^2(\mathbb{R}^3)$ (it is not even a function). This anticipates problems with incorporating some physically plausible states (like the localized states in coordinate or momentum space) into the mathematical formalism of quantum theory

◀ **Historical remark**

- 1800-10: Thomas Young formulates the superposition principle for waves
- 1924: Louis de Broglie introduces the concept of particle wavefunction
- 1926: Erwin Schrödinger formulates wave mechanics
- 1926: Max Born provides the probabilistic interpretation of wavefunction
- 1926-32: John von Neumann formulates QM through linear vector spaces
- 1927-30: Paul Dirac includes into the formulation the δ -function

■ **Quantum measurement**

To explain the which-path version of the double-slit experiment, we assume that the measurement has a dramatic effect on a quantum system: “**reduction**” or “**collapse**” of its wavefunction to the single alternative that was observed.

► **Change of wavefunction in measurement**


Example: position measurement detecting the particle (in time t_0) within the box $(x'_1 \pm \frac{\epsilon_1}{2}, x'_2 \pm \frac{\epsilon_2}{2}, x'_3 \pm \frac{\epsilon_3}{2}) \Rightarrow$ the wavefunction changes as:

$$\psi(\vec{x}, t_0) \text{ delocalized} \xrightarrow{\text{reduction}} \psi(\vec{x}, t_0 + dt) \propto \delta_\epsilon(\vec{x} - \vec{x}') \psi(\vec{x}, t_0) \text{ localized}$$

In an *ideal* ($\epsilon \rightarrow 0$) measurement that detects the particle at \vec{x}' :

$$\psi(\vec{x}, t) \xrightarrow{\text{reduction}} \delta(\vec{x} - \vec{x}')$$

After the position measurement, the wavefunction evolves from a localized one



► Description of the which-path setup in the double-slit experiment

At the double-slit plate:

$$\psi(\vec{x}, t_0) \approx \alpha \delta_A(\vec{x} - \vec{x}_A) + \beta \delta_B(\vec{x} - \vec{x}_B)$$

After which-path measurement ($\delta t \ll \Delta t$):


$$\psi(\vec{x}, t_0 + \delta t) \approx \begin{cases} \delta_A(\vec{x} - \vec{x}_A) & \text{probability} \approx |\alpha|^2 \\ \delta_B(\vec{x} - \vec{x}_B) & \text{probability} \approx |\beta|^2 \end{cases}$$

At the screen:

$$\psi(\vec{x}, t_0 + \Delta t) \approx \begin{cases} \psi_A(\vec{x}, \Delta t) & \text{probability} \approx |\alpha|^2 \\ \psi_B(\vec{x}, \Delta t) & \text{probability} \approx |\beta|^2 \end{cases}$$

$$\Rightarrow \boxed{\rho(\vec{x}) \approx |\alpha \psi_A(\vec{x}, \Delta t)|^2 + |\beta \psi_B(\vec{x}, \Delta t)|^2} \text{ probability distribution on the screen}$$

So the interference pattern is destroyed! This is a direct consequence of the wavefunction collapse caused by the which-path measurement.



Note: Disappearance of the interference pattern can be also induced by the presence of an additional quantum system (an “atom”) that interacts with the particle inside the two-slit device so that it records the which-path information—without any observer actually reading it! The composite particle–atom system is described by an extended wavefunction with both particle & atom degrees of freedom. The measurement-like effect then follows from a continuous, collapse-free evolution of the extended wavefunction reflecting the particle–atom interaction. The collapse assumption is nevertheless useful if we want to describe the measured system autonomously, irrespective of the “measuring agents”.

► **Summing amplitudes versus summing probabilities:** For a general branching processes with disjunct alternative paths A & B (real or symbolic), the probability to pass the branching while the path is not *explicitly* measured depends on whether the paths can/cannot, *in principle*, be distinguished:

For **indistinguishable paths** we sum amplitudes: $\mathbf{a} \propto \mathbf{a}_A + \mathbf{a}_B$

\Rightarrow interference effects occur in $\mathbf{p} = |\mathbf{a}|^2$

For **distinguishable paths** we sum probabilities: $\mathbf{p} \propto \mathbf{p}_A + \mathbf{p}_B$

\Rightarrow interference effects do not occur

► **Quantum logic:** An attempt was made to assign the strange properties of the quantum world to a non-classical underlying logic. In the double-slit experiment it can be introduced via the following “propositions”:

$A, B \equiv$ passage through slit A,B $S \equiv$ detection at given place of screen
 Different outcomes of interference & which-path setups indicate that:

$$\boxed{\underbrace{(A \vee B) \wedge S}_{\text{interference setup}} \neq \underbrace{(A \wedge S) \vee (B \wedge S)}_{\text{which-path setup}}} \quad (\text{where } \vee \equiv \text{“or” and } \wedge \equiv \text{“and”}) \\ \Rightarrow \text{violation of a common logic axiom}$$

◀ Historical remark

1924-35: Bohr (Copenhagen) versus Einstein debate. Niels Bohr defends a “subjective” approach (with the observer playing a role in the “creation” of reality)

1927: the first explicit note of wavefunction collapse by Werner Heisenberg

1932: inclusion of collapse into the mathematical formulation of QM by John von Neumann (discussions about its physical meaning continue up to now)

1936: Garrett Birkhoff and J. von Neumann formally introduce quantum logic

1a. SPACE OF QUANTUM STATES

Quantum theory has rather sophisticated formalism based on the mathematics matured at the beginning of the 20th century. Its interpretation in terms of “common sense” becomes a nontrivial issue rising questions about the link of physical theory to reality. The problem starts already on the deepest level—with the definition of states of quantum systems, i.e., sets of attributes sufficient for a unique description of the system’s evolution. While the mathematical representation of states in classical physics is rather intuitive and comprehensible (using the notion of phase space), quantum physics resorts to much more abstract ideas.

Roughly the first half of this book attempts to give a complete overview of the quantum formalism. The chapters that contain letter “a” in the numbering outline, step by step, the basic elements of the mathematical description. The chapters with letter “b” give some simple concrete examples (mostly in single-particle systems) of the respective ideas. To keep immediate link between the *Geist* and *Substanz*, we present the “a” and “b” chapters in an alternating, entangled way.


■ Hilbert space

To capture the quantum uncertainty, i.e., the possibility of different outcomes of various measurements performed on systems in the same state, we will assume that *distinct states of the system are not always perfectly distinguishable*. The states may show some “overlaps”, which allow one to identify a given state with another state—e.g., the state of a particle described by a delocalized wavefunction $\psi(\vec{x})$ with a state localized at a single place \vec{x}' . This means that the states are not represented by isolated points *à la* points in the classical phase space. Instead, they can be associated with **vectors** in linear vector spaces. If two vectors are not perpendicular to each other, they have a common component

whose size sets limits to their mutual distinguishability.

► State of a physical system

The state represents a complete set of parameters characterizing a physical system in the sense of an autonomous determinism: The knowledge of state at a *single time* ($t=0$) suffices to determine the state at *any time* in past or future ($t \geq 0$). Let $|\psi\rangle$ denote a mathematical entity describing an arbitrary physical state of a given quantum system (short-cut: $|\psi\rangle \equiv$ “a state”). Let \mathcal{H} be a system-specific space containing all such entities (**state space**). We make our first fundamental assumption:



The space of states \mathcal{H} of an arbitrary quantum system is a **Hilbert space**, in which individual states are represented by rays of vectors.

The Hilbert space is defined by the following 3 requirements:

► Requirement 1: The space \mathcal{H} supports the **superposition principle**

$$\left. \begin{array}{l} |\psi_1\rangle, |\psi_2\rangle \in \mathcal{H} \\ \alpha, \beta \in \mathbb{C} \end{array} \right\} \Rightarrow \boxed{|\psi\rangle = \alpha|\psi_1\rangle + \beta|\psi_2\rangle \in \mathcal{H}} \quad \begin{array}{l} \text{superposition of states} \\ |\psi_1\rangle \text{ and } |\psi_2\rangle \end{array}$$

$\Rightarrow \mathcal{H}$ is a **complex vector space**

Why we need superpositions: To describe the single-particle interference in the double-slit experiment (Intro.), we must add the waves from both slits.

► Requirement 2: The space \mathcal{H} supports a **scalar product** $\langle \psi_1 | \psi_2 \rangle \in \mathbb{C}$

Properties: $\langle \psi_1 | \psi_2 \rangle = \langle \psi_2 | \psi_1 \rangle^*$, $\langle \psi_1 | \alpha\psi_2 + \beta\psi_3 \rangle = \alpha\langle \psi_1 | \psi_2 \rangle + \beta\langle \psi_1 | \psi_3 \rangle$, $\langle \psi | \psi \rangle \geq 0$

Normalization of state vectors: Real number $\boxed{\|\psi\| \equiv \sqrt{\langle \psi | \psi \rangle} \geq 0}$ is a norm of $|\psi\rangle$. Scaling of state vectors, i.e. multiplication $|\psi'\rangle = a|\psi\rangle$ by any constant $a \in \mathbb{C}$, does not change their physical content (so both $|\psi'\rangle$, $|\psi\rangle$ describe the same state). Hence any state vector can be scaled so that it becomes normalized:

$$\boxed{\langle \psi | \psi \rangle = 1}$$

In QM we use normalized vectors, but this cannot be set as a constraint in \mathcal{H} because of the superposition principle (if linearly combining two normalized states, the resulting superposition is generally not normalized).


Distance of 2 vectors: $d^2(\psi_1, \psi_2) \equiv \|\psi_1 - \psi_2\|^2 = \langle \psi_1 | \psi_1 \rangle + \langle \psi_2 | \psi_2 \rangle - 2\text{Re}\langle \psi_1 | \psi_2 \rangle$

Schwarz inequality for normalized vectors: $\boxed{|\langle \psi_1 | \psi_2 \rangle|^2 \leq \underbrace{\langle \psi_1 | \psi_1 \rangle}_1 \underbrace{\langle \psi_2 | \psi_2 \rangle}_1 = 1}$

Why we need scalar product:

Results of quantum measurements are generally indeterministic (described in the probabilistic way, see Intro. & Sec. 2a.). A single measurement does not allow one to uniquely determine the state. The possibility to identify state $|\psi_2\rangle$

with $|\psi_1\rangle$ or vice versa in an “optimal” single measurement is determined by the overlap of the corresponding vectors. For $\|\psi_1\|=\|\psi_2\|=1$ we have:



$$\underbrace{\mathbf{a}_{\psi_2}(\psi_1) \equiv \langle \psi_1 | \psi_2 \rangle}_{\text{amplitude}} \quad \underbrace{\mathbf{p}_{\psi_2}(\psi_1) \equiv |\langle \psi_1 | \psi_2 \rangle|^2}_{\text{probability}}$$

Number $\mathbf{a}_{\psi_2}(\psi_1) \in \mathbb{C}$ satisfying $|\mathbf{a}_{\psi_2}(\psi_1)| \in [0, 1]$ represents **amplitude** for finding $|\psi_1\rangle$ in $|\psi_2\rangle$

The corresponding **probability** $\mathbf{p}_{\psi_2}(\psi_1) \in [0, 1]$ is obtained by squaring the amplitude’s modulus

Consequence:

States $|\psi_1\rangle, |\psi_2\rangle$ are perfectly **distinguishable** iff orthogonal: $\langle \psi_1 | \psi_2 \rangle = 0$

► **Requirement 3:** \mathcal{H} is **complete**, i.e. \forall converging sequence (in the Cauchy sense with distance d) of vectors $\{|\psi_i\rangle\}_i$ the limit $\lim_{i \rightarrow \infty} |\psi_i\rangle \equiv |\psi_\infty\rangle \in \mathcal{H}$. This shall avoid problems with missing limits (unfortunately, it does not apply to the δ -function, see Intro., as the “convergence” to δ is not of the Cauchy type).

► Separable Hilbert spaces

\mathcal{H} is separable if it has a **countable** (possibly finite) **set of basis vectors**

We can choose an **orthonormal basis** $\{|\phi_i\rangle\}_{i=1}^{d_{\mathcal{H}}}$ satisfying $\langle \phi_i | \phi_j \rangle = \delta_{ij}$

The number of basis vectors $d_{\mathcal{H}}$ is called dimension of \mathcal{H}

⇒ Each state $|\psi\rangle$ can be expressed as a unique complex superposition of basis vectors:

Normalization:

$$\langle \psi | \psi \rangle = \sum_{i=1}^{d_{\mathcal{H}}} \sum_{j=1}^{d_{\mathcal{H}}} \alpha_i^* \alpha_j \overbrace{\langle \phi_i | \phi_j \rangle}^{\delta_{ij}} = \sum_{i=1}^{d_{\mathcal{H}}} |\alpha_i|^2 = 1$$

$$|\psi\rangle = \sum_{i=1}^{d_{\mathcal{H}}} \underbrace{\langle \phi_i | \psi \rangle}_{\alpha_i} |\phi_i\rangle$$

Applicability: Systems with finite numbers of particles, systems with finite numbers of degrees of freedom (possibly selected subsets of degrees of freedom)

Isomorphism of separable Hilbert spaces

Any separable \mathcal{H} with an infinite basis set is isomorphic with the **space** ℓ^2 formed by infinite “columns” of complex numbers $\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \end{pmatrix}$ satisfying $\sum_{i=1}^{\infty} |\alpha_i|^2 < \infty$

Mapping $\mathcal{H} \rightarrow \ell^2$: Expansion coefficients $\langle \phi_i | \psi \rangle$ of a chosen vector $|\psi\rangle \in \mathcal{H}$ in a given basis $\{|\phi_i\rangle\}_i$ are associated with the numbers α_i defining the vector $\in \ell^2$

Superpositions $a|\psi\rangle + b|\psi'\rangle$ mapped onto: $\begin{pmatrix} a\alpha_1 + b\alpha'_1 \\ a\alpha_2 + b\alpha'_2 \\ \vdots \end{pmatrix}$

Scalar product represented by: $\langle \psi | \psi' \rangle \equiv \sum_i \alpha_i^* \alpha'_i = (\alpha_1^*, \alpha_2^*, \dots) \begin{pmatrix} \alpha'_1 \\ \alpha'_2 \\ \vdots \end{pmatrix}$

► Nonseparable Hilbert spaces

\mathcal{H} is nonseparable if it has no countable basis. This applies in systems with unbounded particle numbers, quantum fields, continuum...

◀ Historical remark

1900-10: David Hilbert (with E. Schmidt) introduces the ∞ -dimensional space of square-integrable functions and elaborates the theory of such spaces

1927: John von Neumann (working under Hilbert) introduces abstract Hilbert spaces into QM (1932: book *Mathematische Grundlagen der Quantenmechanik*)

■ Rigged Hilbert space

Although the standard Hilbert space is sufficient for consistent formulation of QM, we will see soon that its suitable extension is very helpful.

► Hierarchy of spaces based on $\mathcal{H} \equiv \ell^2$

$\underline{\mathcal{H}}$ is a space of sequences $\{\alpha_i\}_{i=1}^{\infty} \equiv |\psi\rangle$ satisfying $\sum_i |\alpha_i|^2 i^m < \infty$ for $m=0,1,2,\dots$

These form a dense subset of ℓ^2

$\overline{\mathcal{H}}$ (conjugate space to $\underline{\mathcal{H}}$) is a space of sequences $\{\alpha'_i\}_{i=1}^{\infty} \equiv |\psi'\rangle$ satisfying

$\langle \psi | \psi' \rangle < \infty$ for any $|\psi\rangle \in \underline{\mathcal{H}}$. This set contains ℓ^2 as a subset.


$\langle \psi | \psi' \rangle \equiv \sum_i \alpha_i^* \alpha'_i < \infty \Rightarrow \sum_i |\alpha'_i|^2 \frac{1}{i^m} < \infty \Rightarrow |\alpha'_i|^2$ may polynomially diverge

In general, the smaller is $\underline{\mathcal{H}}$, the larger is $\overline{\mathcal{H}}$

$\underline{\mathcal{H}}$ and $\overline{\mathcal{H}}$ are linear vector spaces, but not Hilbert spaces:

$\underline{\mathcal{H}}$ is not complete

$\overline{\mathcal{H}}$ does not have scalar product



► Gelfand triple $\underline{\mathcal{H}} \subset \mathcal{H} \subset \overline{\mathcal{H}}$

This “sandwich” of spaces is sometimes called the “rigged Hilbert space”, indicating that only such an extended structure allows one to “safely sail the sea” of quantum physics. It turns out that solutions of some basic quantum problems is out of \mathcal{H} but belongs to the larger space $\overline{\mathcal{H}}$, while the definition domain of some quantum operators is not \mathcal{H} but rather its subspace $\underline{\mathcal{H}}$ (see Secs. 2a & 2b).

■ Dirac notation

Physicists are proud to master a symbolic technique that makes some involved mathematical reductions much easier to follow. Although the “bra-ket” formalism is not always fully rigorous, it is extremely efficient especially when dealing with the action of linear operators in Hilbert spaces.

► Kets and bras

For any vector $|\psi\rangle \in \mathcal{H}$, called **ket**, there exists a linear functional $F_\psi \equiv \langle \psi|$,

called **bra**, such that the value assigned by F_ψ to $|\phi\rangle \in \mathcal{H}$ is $F_\psi(\phi) \equiv \langle\psi|\phi\rangle$ (the words following from “bra-c-ket”)


The bras also satisfy the superposition principle:

$$\alpha\langle\psi_1| + \beta\langle\psi_2| \equiv \langle\alpha^*\psi_1 + \beta^*\psi_2|$$

and the spaces of kets & bras are isomorphic.

Matrix forms:

$$\langle\psi| \equiv (\alpha_1^*, \alpha_2^*, \dots) \quad \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \end{pmatrix} \equiv |\psi\rangle$$



► Linear operators

Linear operators play a very important role in QM. They will be subject to systematic study from Sec. 2a. Here we just introduce basic notions.

Linear operator $\hat{O}|\psi\rangle = |\psi'\rangle$ is a mapping $\mathcal{H} \rightarrow \mathcal{H}$ of the Hilbert space to itself satisfying the **linearity condition**: $\hat{O}(\alpha|\psi_1\rangle + \beta|\psi_2\rangle) = \alpha\hat{O}|\psi_1\rangle + \beta\hat{O}|\psi_2\rangle$

$\Rightarrow \hat{O}$ is completely defined via its action on any basis: $\{|\phi_i\rangle\}_{i=1}^{d_{\mathcal{H}}} \xrightarrow{\hat{O}} \{|\phi'_i\rangle\}_{i=1}^{d_{\mathcal{H}}}$

$$\Rightarrow \hat{O}|\psi\rangle = \sum_i \underbrace{\langle\phi_i|\psi\rangle}_{|\phi_i\rangle} \underbrace{\hat{O}|\phi_i\rangle}_{|\phi'_i\rangle} \Rightarrow \hat{O} \equiv \sum_{i=1}^{d_{\mathcal{H}}} |\phi'_i\rangle\langle\phi_i| \quad \begin{matrix} |\phi'_i\rangle\langle\phi_i| \\ \Downarrow \\ |\phi_i\rangle \xrightarrow{\hat{O}} |\phi'_i\rangle \end{matrix}$$

Any expression of the form $|\phi'\rangle\langle\phi|$ is a linear operator: $|\psi\rangle \xrightarrow{\hat{O}} \langle\phi|\psi\rangle|\phi'\rangle$.

Any linear operator \hat{O} can be expressed as a sum over terms $\propto |\phi_j\rangle\langle\phi_i|$ containing vectors of the *same* basis. This is achieved via the identity (unit) operator:

$$\sum_{i=1}^{d_{\mathcal{H}}} |\phi_i\rangle\langle\phi_i| = \hat{I} \quad \{|\phi_i\rangle\} \xrightarrow{\hat{I}} \{|\phi_i\rangle\} \quad \hat{O} \equiv \text{unit operator} \Rightarrow \hat{O} = \sum_{i=1}^{d_{\mathcal{H}}} \sum_{j=1}^{d_{\mathcal{H}}} \underbrace{\langle\phi_j|\hat{O}\phi_i\rangle}_{\langle\phi_j|\hat{O}\phi_i\rangle \equiv O_{ji}} |\phi_j\rangle\langle\phi_i|$$

Matrix form
general linear operator: $\hat{O} \equiv \begin{pmatrix} O_{11} & O_{12} & \dots \\ O_{21} & O_{22} & \\ \vdots & & \ddots \end{pmatrix}$

► Projectors


Projection operators (projectors) are linear operators satisfying $\hat{P}^2 = \hat{P}$ (i.e., repeated projection is redundant)

Let $\{|\phi_i\rangle\}_{i=1}^{d_0} \equiv$ orthonormal basis of a subspace $\mathcal{H}_0 \subset \mathcal{H}$. We have $\langle\phi_i|\phi_j\rangle = \delta_{ij}$

$$\hat{P}_0 = \sum_{i=1}^{d_0} |\phi_i\rangle\langle\phi_i| \quad \text{is a projector to } \mathcal{H}_0: \quad \hat{P}_0|\psi\rangle \begin{cases} = 0 & \text{for } |\psi\rangle \perp \mathcal{H}_0 \\ \in \mathcal{H}_0 & \text{otherwise} \end{cases}$$

Completeness relation: the projector to the whole \mathcal{H} is the identity operator (see above):

$$\hat{P}_{\mathcal{H}} = \sum_{i=1}^{d_{\mathcal{H}}} |\phi_i\rangle\langle\phi_i| = \hat{I}$$



Probability to identify $|\psi\rangle$ with *any* state from the subspace \mathcal{H}_0 :

In generalization of the above formula $\mathbf{p}_\psi(\psi_0) = |\langle\psi_0|\psi\rangle|^2 = \langle\psi|\psi_0\rangle\langle\psi_0|\psi\rangle$, the overall probability to (incorrectly)

associate a given state $|\psi\rangle \in \mathcal{H}$ with an arbitrary state $|\psi'\rangle \in \mathcal{H}_0$ is given by:

$$\mathbf{p}_\psi(\mathcal{H}_0) \equiv \langle\psi|\hat{P}_0|\psi\rangle = \sum_{i=1}^{d_0} |\langle\phi_i|\psi\rangle|^2$$

Matrix form of projector operators:

In an orthonormal basis $\{|\phi_i\rangle\}_{i=1}^{d_{\mathcal{H}}}$ of \mathcal{H} containing as a subset the basis $\{|\phi_{i_j}\rangle\}_{j=1}^{d_0}$ of \mathcal{H}_0 (with $i_j \equiv$ indices of the \mathcal{H}_0 basis vectors in the \mathcal{H} basis), the projector is expressed as a diagonal matrix with d_0 units and $(d_{\mathcal{H}}-d_0)$ zeros on the diagonal:

$$\hat{P}_0 = \begin{pmatrix} X_1 & 0 & \dots \\ 0 & X_2 & \dots \\ \vdots & & \ddots \end{pmatrix} \text{ with } X_i = \begin{cases} 1 & \text{for } i \in \{i_1, i_2, \dots, i_{d_0}\} \\ 0 & \text{for } i \notin \{i_1, i_2, \dots, i_{d_0}\} \end{cases}$$

◀ Historical remark

1930: Paul Dirac writes the book *The Principles of Quantum Mechanics*, which provides a more intuitive (compared to von Neumann) path to quantum theory, using non-normalizable vectors and δ -function (bra-kets in 3rd edition 1947)

1950-60's: I.M. Gelfand & N.Y. Vilenkin introduce rigged Hilbert spaces, putting Dirac's approach on more rigorous grounds. Systematic use in QM since 1966 (by A. Böhm et al.) but up to now rather scarce

■ Summing Hilbert spaces

One can combine one or more Hilbert spaces in the style of summation. The resulting space then contains the summed spaces as ordinary subspaces.

► Direct sum

Let $\{|\phi_{1i}\rangle\}_{i=1}^{d_1}$ be an orthonormal basis of \mathcal{H}_1 and $\{|\phi_{2j}\rangle\}_{j=1}^{d_2}$ one of \mathcal{H}_2

Direct sum space $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ has the "summed" basis $|\Phi_{ki}\rangle \equiv \begin{cases} |\phi_{1i}\rangle & \text{for } k=1 \\ |\phi_{2i}\rangle & \text{for } k=2 \end{cases}$

Hence $\mathcal{H}_1 \oplus \mathcal{H}_2$ consists of all normalizable linear combinations of the basis vectors $|\Phi_{ki}\rangle$ formed by a unification of the basis vectors of \mathcal{H}_1 and \mathcal{H}_2 .

Dimension: $d_{\mathcal{H}_1 \oplus \mathcal{H}_2} = d_1 + d_2$ Orthonormality of basis: $\langle\Phi_{ki}|\Phi_{k'i'}\rangle = \delta_{kk'}\delta_{ii'}$

► State decomposition

Any vector $|\Psi\rangle = \sum_{k,i} \alpha_{ki} |\Phi_{ki}\rangle \in \mathcal{H}$ can be written as $|\Psi\rangle = |\psi_1\rangle + |\psi_2\rangle$ with $|\psi_k\rangle \in \mathcal{H}_k$ ($k=1, 2$)

$$|\Psi\rangle = \underbrace{\sum_{i=1}^{d_1} \alpha_{1i} |\phi_{1i}\rangle}_{|\psi_1\rangle \equiv \hat{P}_1|\Psi\rangle \in \mathcal{H}_1} + \underbrace{\sum_{j=1}^{d_2} \alpha_{2j} |\phi_{2j}\rangle}_{|\psi_2\rangle \equiv \hat{P}_2|\Psi\rangle \in \mathcal{H}_2}$$

Projectors to the subspaces \mathcal{H}_k

$\hat{P}_k = \sum_{i=1}^{d_k} |\Phi_{ki}\rangle\langle\Phi_{ki}| \Rightarrow \begin{cases} \text{orthogonality : } \hat{P}_1\hat{P}_2 = \hat{P}_2\hat{P}_1 = 0 \\ \text{completeness : } \hat{P}_1 + \hat{P}_2 = \hat{I}_{\mathcal{H}} \end{cases}$

Scalar product: $\langle\Psi|\Psi'\rangle_{\mathcal{H}} = \langle\psi_1|\psi'_1\rangle_{\mathcal{H}_1} + \langle\psi_2|\psi'_2\rangle_{\mathcal{H}_2}$ where $\begin{cases} |\psi_k\rangle = \hat{P}_k|\Psi\rangle \\ |\psi'_k\rangle = \hat{P}_k|\Psi'\rangle \end{cases}$

► Finite-dimensional matrix representation:

$$|\psi_1\rangle = \begin{pmatrix} \alpha_{11} \\ \vdots \\ \alpha_{1d_1} \end{pmatrix}, |\psi_2\rangle = \begin{pmatrix} \alpha_{21} \\ \vdots \\ \alpha_{2d_2} \end{pmatrix} \Rightarrow |\Psi\rangle = \begin{pmatrix} \alpha_{11} \\ \vdots \\ \alpha_{1d_1} \\ \dots \\ \alpha_{21} \\ \vdots \\ \alpha_{2d_2} \end{pmatrix}$$

► Multiple sums of Hilbert spaces: $\mathcal{H} = \bigoplus_{k=1}^n \mathcal{H}_k$

► **The use in QM:** $\mathcal{H}_k \subset \mathcal{H}$ can be associated with subspaces corresponding to various values a_k of an observable A (states $|\psi\rangle \in \mathcal{H}_k$ yield output a_k with certainty; see Sec. 2a). Subspaces in the direct sum can also collect vectors with different symmetry properties (e.g., subspaces of even and odd wavefunctions).

■ Multiplying Hilbert spaces

Hilbert spaces can also be combined in the style of multiplication. This commonly happens in **composite quantum systems** that consist of two or more subsystems (several particles or distinct subsets of degrees of freedom). The multiplication is a rather interesting operation since it allows one to create so called *entangled quantum states* which have no analogue in the classical world.

► **Direct (tensor) product**

Let $\{|\phi_{1i}\rangle\}_{i=1}^{d_1}$ be an orthonormal basis of \mathcal{H}_1 and $\{|\phi_{2j}\rangle\}_{j=1}^{d_2}$ one of \mathcal{H}_2

Tensor product space $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$ has the product basis $|\Phi_{ij}\rangle \equiv |\phi_{1i}\rangle|\phi_{2j}\rangle$

This means that $\mathcal{H}_1 \otimes \mathcal{H}_2$ consists of all normalizable linear combinations of the basis vectors $|\Phi_{ij}\rangle$ formed by direct products of \mathcal{H}_1 and \mathcal{H}_2 basis vectors. Note that non-product bases of $\mathcal{H}_1 \otimes \mathcal{H}_2$ can also be constructed.

Dimension: $d_{\mathcal{H}_1 \otimes \mathcal{H}_2} = d_1 \cdot d_2$ Orthonormality of basis: $\langle \Phi_{ij} | \Phi_{i'j'} \rangle = \delta_{ii'} \delta_{jj'}$

► **Factorized states**

For any pair of states $\left\{ \begin{array}{l} |\psi_1\rangle = \sum_i \alpha_i |\phi_{1i}\rangle \in \mathcal{H}_1 \\ |\psi_2\rangle = \sum_j \beta_j |\phi_{2j}\rangle \in \mathcal{H}_2 \end{array} \right\}$ there exists the product state

$$|\Psi_{\otimes}\rangle \equiv \underbrace{|\psi_1\rangle \otimes |\psi_2\rangle}_{\equiv |\psi_1\rangle_1 |\psi_2\rangle_2} = \sum_{i=1}^{d_1} \sum_{j=1}^{d_2} \underbrace{\alpha_i \beta_j}_{\gamma_{ij}} |\Phi_{ij}\rangle$$

A factorized state allows one to uniquely identify the associated state vectors of individual subsystems

Scalar product for factorized states: $\langle \Psi_{\otimes} | \Psi'_{\otimes} \rangle_{\mathcal{H}} = \langle \psi_1 | \psi'_1 \rangle_{\mathcal{H}_1} \cdot \langle \psi_2 | \psi'_2 \rangle_{\mathcal{H}_2}$

► **Entangled states**

The possibility to express coefficients γ_{ij} of a general superposition $\sum_{ij} \gamma_{ij} |\Phi_{ij}\rangle$ in the above factorized form ($\gamma_{ij} = \alpha_i \beta_j$) is rather scarce. Almost all states in $\mathcal{H}_1 \otimes \mathcal{H}_2$ are unfactorizable, so called entangled states:

$$|\Psi\rangle = \sum_{i=1}^{d_1} \sum_{j=1}^{d_2} \underbrace{\gamma_{ij}}_{\neq \alpha_i \beta_j} |\Phi_{ij}\rangle \neq |\psi_1\rangle_1 |\psi_2\rangle_2$$

An entangled state does *not* have any associated state vectors of individual subsystems

► Multiple products of Hilbert spaces:

$$\mathcal{H} = \bigotimes_{k=1}^n \mathcal{H}_k$$

More and less precise notations:

$$\mathcal{H}_k \ni |\psi\rangle \equiv |\psi\rangle_k \quad \text{and} \quad \bigotimes_{k=1}^n \mathcal{H}_k \ni |\psi_1\rangle_1 \otimes |\psi_2\rangle_2 \dots \otimes |\psi_n\rangle_n \equiv |\psi_1\rangle_1 |\psi_2\rangle_2 \dots |\psi_n\rangle_n$$

► **The use in QM**

Hilbert space \mathcal{H} of a composite system is the \otimes product of partial spaces \mathcal{H}_k . The multiplied spaces \mathcal{H}_k can be associated with the spaces corresponding to different parts of the system (e.g. different particles) or to different dynamical variables (e.g., spatial and spin degrees of freedom). Entangled state vectors correspond to *non-classical* situations in which only the whole system and not its individual parts are attributed by a pure quantum-mechanical state (the subsystems are in so called mixed states, see Sec. 6a). Entanglement represents a genuinely **quantum correlation** of the system's parts.

◀ Historical remark

1935: A. Einstein, B. Podolsky & N. Rosen use an entangled state to claim that QM is incomplete. E. Schrödinger analyzes such states and coins the term “entanglement”

1b. EXAMPLES OF QUANTUM HILBERT SPACES

In the following, we describe specific state spaces for particles with spin 0 and $\frac{1}{2}$, and the spaces assigned to collections of such particles. We meet another essentially quantum phenomenon: indistinguishability of particles. And we introduce the space of qubits — the playground of quantum information technologies.

■ Single structureless and spinless particle

Particles with no internal degrees of freedom are described by ordinary scalar wavefunctions (cf. Intro.).

► **Wavefunction** $\psi(\vec{x}) \equiv |\psi\rangle \in$

$$\mathcal{H} \equiv \mathcal{L}^2(\mathbb{R}^3)$$

Hilbert space of **square-integrable functions**

$$\langle \psi_1 | \psi_2 \rangle \equiv \int \psi_1^*(\vec{x}) \psi_2(\vec{x}) d\vec{x} \equiv \int \psi_1^*(\vec{y}) \psi_2(\vec{y}) \left| \text{Det} \frac{\partial(x_1 \dots x_3)}{\partial(y_1 \dots y_3)} \right| d\vec{y} \quad \text{scalar product}$$


Cartesian & curvilinear coordinates

Expansion of $\psi(\vec{x})$ in a discrete **basis of orthonormal functions** $\{\phi_i(\vec{x})\}_{i=1}^{\infty}$
 \Rightarrow isomorphism of $\mathcal{L}^2(\mathbb{R}^3)$ with ℓ^2

Normalization: $\langle \psi | \psi \rangle = \int |\psi(\vec{x})|^2 d\vec{x} = 1$

Probabilistic interpretation:

$|\psi(\vec{x})|^2 \equiv \rho(\vec{x})$ is the probability density for finding the particle at position \vec{x} . This follows from the association of the state $|\vec{x}'\rangle$ of the particle at a single sharp position \vec{x}' with the



Dirac δ -function $\delta(\vec{x}-\vec{x}')$ (see Sec. 0). Hence the amplitude & probability densities read:

$$\begin{aligned} \mathbf{a}_\psi(\vec{x}') &= \langle \vec{x}' | \psi \rangle = \int \delta(\vec{x}-\vec{x}') \psi(\vec{x}) d\vec{x} = \psi(\vec{x}') \\ \mathbf{p}_\psi(\vec{x}') &= |\mathbf{a}_\psi(\vec{x}')|^2 = |\psi(\vec{x}')|^2 \equiv \rho(\vec{x}') \end{aligned}$$

Probability expressions for other observables will be treated in Secs. 2a & 2b.

► Rigged Hilbert space of wavefunctions

There is a problem that localized states $|\vec{x}'\rangle \equiv \delta(\vec{x}-\vec{x}')$ as well as other important states (like plane waves $e^{i\vec{k}\cdot\vec{x}}$, see Sec. 2b) are not in $\mathcal{L}^2(\mathbb{R}^3)$ (they are not quadratically integrable). The rescue comes with the introduction of a convenient Gelfand triple $\underline{\mathcal{H}} \subset \mathcal{H} \subset \overline{\mathcal{H}}$ of spaces. In the 1D case, we define:

$\underline{\mathcal{H}} \equiv$ dense subset of functions: $\int_{-\infty}^{+\infty} |\psi(x)|^2 (1+|x|)^m dx < \infty$ for $m = 0, 1, 2, \dots$

$\overline{\mathcal{H}} \equiv$ functions satisfying $\int_{-\infty}^{+\infty} \psi'^* \psi dx < \infty \forall \psi' \in \underline{\mathcal{H}}$

Then $\overline{\mathcal{H}}$ includes also polynomially diverging functions, plane waves, δ -functions

An alternative mathematically consistent approach (see Sec. 2a) is to consider only imperfectly localized states, like those within interval $x_i \in [x'_i - \frac{\epsilon_i}{2}, x'_i + \frac{\epsilon_i}{2}]$ around \vec{x}' represented by wavefunctions $\delta_{\vec{\epsilon}}(\vec{x}-\vec{x}') \in \mathcal{L}^2(\mathbb{R}^3)$.

◀ Historical remark

1926: Erwin Schrödinger formulates QM in terms of wavefunction and Max Born develops its probabilistic interpretation

■ Single structureless particle with spin $\frac{1}{2}$

Electrons have spin $\frac{1}{2}$. The spin is a genuinely quantum feature of a particle, which (even for point-like elementary particles) is understood as an intrinsic unstopable rotation. The general theory of angular momentum in QM will be developed in Secs. 3b & 4b, here we just introduce spinor wavefunctions as the simplest two-component generalization of scalar wavefunctions.

► Spin = intrinsic angular momentum of a particle

The lowest nonzero spin is $s = \frac{1}{2}$, having only 2 possible projections (spin states) $s_\bullet = \pm s\hbar$ in an arbitrarily chosen spatial direction \bullet (conventionally $\bullet = z$):

$$\left. \begin{array}{l} \text{spin up} \quad s_z = +\frac{\hbar}{2} \Rightarrow |\uparrow\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix} \equiv |\chi_1\rangle \\ \text{spin down} \quad s_z = -\frac{\hbar}{2} \Rightarrow |\downarrow\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix} \equiv |\chi_2\rangle \end{array} \right\} \Rightarrow \text{general state } |\psi\rangle = \alpha_1 |\uparrow\rangle + \alpha_2 |\downarrow\rangle \equiv \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}$$

► Spin Hilbert space: $\mathcal{H} \equiv \mathbb{C}^2$ with $\langle \psi | \psi' \rangle \equiv (\alpha_1^*, \alpha_2^*) \begin{pmatrix} \alpha_1' \\ \alpha_2' \end{pmatrix} = \alpha_1^* \alpha_1' + \alpha_2^* \alpha_2'$

Normalization: $|\alpha_1|^2 + |\alpha_2|^2 = 1$

Probability to find the spin up/down:

$$\begin{aligned} \mathbf{p}_\psi(\uparrow) &= |\mathbf{a}_\psi(\uparrow)|^2 = |\langle \uparrow | \psi \rangle|^2 = |\alpha_1|^2 \\ \mathbf{p}_\psi(\downarrow) &= |\mathbf{a}_\psi(\downarrow)|^2 = |\langle \downarrow | \psi \rangle|^2 = |\alpha_2|^2 \end{aligned}$$

For probabilities of spin projections in an arbitrary direction see Sec. 2b.

Note: General spin $s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$ has $(2s+1)$ spin projections $s_\bullet = -s\hbar, (-s+1)\hbar, \dots, (+s-1)\hbar, +s\hbar$ to any direction, so $\mathcal{H} \equiv \mathbb{C}^{2s+1}$ (see Sec. 3b).

► **Both spatial and spin degrees of freedom**

⇒ direct product of spatial and spin Hilbert spaces:

$$\mathcal{H} \equiv \mathcal{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^2$$


Basis vectors: $|\Phi_{ij}\rangle = |\phi_i\rangle|\chi_j\rangle$, where $\{|\phi_i\rangle\}_{i=1}^{\infty}$ is an arbitrary basis in $\mathcal{L}^2(\mathbb{R}^3)$

General state expansion: $|\psi\rangle = \sum_{i=1}^{\infty} \sum_{j=1}^2 \alpha_{ij} |\phi_i\rangle|\chi_j\rangle = \sum_{i=1}^{\infty} [\alpha_{i1} \phi_i(\vec{x}) |\uparrow\rangle + \alpha_{i2} \phi_i(\vec{x}) |\downarrow\rangle]$

$$= \sum_{i=1}^{\infty} \begin{pmatrix} \alpha_{i1} \\ \alpha_{i2} \end{pmatrix} \phi_i(\vec{x}) = \begin{pmatrix} \sum_i \alpha_{i1} \phi_i(\vec{x}) \\ \sum_i \alpha_{i2} \phi_i(\vec{x}) \end{pmatrix} = \begin{pmatrix} \psi_{\uparrow}(\vec{x}) \\ \psi_{\downarrow}(\vec{x}) \end{pmatrix} \equiv \boldsymbol{\psi}(\vec{x}) \equiv \psi(\vec{x}, m_s) \quad \text{spinor}$$

$\underbrace{}_{\pm \frac{1}{2}}$

Spinor is a two-component wavefunction equivalent to a wavefunction with a continuous variable \vec{x} and a discrete two-valued variable m_s (the spin projection to z -direction). Note that transformation properties of spinors under spatial rotations are different from ordinary vectors (they will be derived in Sec. 4b).



Simplified notation: $(\vec{x}, m_s) \equiv \boldsymbol{\xi}$ with $\int d\boldsymbol{\xi} \equiv \sum_{m_s} \int d\vec{x}$

► **Scalar product** $\langle \psi | \psi' \rangle = \sum_{ij} \sum_{i'j'} \alpha_{ij}^* \alpha'_{i'j'} \underbrace{\langle \phi_i | \phi_{i'} \rangle}_{\delta_{ii'}} \underbrace{\langle \chi_j | \chi_{j'} \rangle}_{\delta_{jj'}} = \sum_{ij} \alpha_{ij}^* \alpha'_{ij}$
can be expressed as:

$$\langle \psi | \psi' \rangle \equiv \int (\psi_{\uparrow}^*(\vec{x}), \psi_{\downarrow}^*(\vec{x})) \begin{pmatrix} \psi'_{\uparrow}(\vec{x}) \\ \psi'_{\downarrow}(\vec{x}) \end{pmatrix} d\vec{x} = \sum_{m_s} \int \psi^*(\vec{x}, m_s) \psi'(\vec{x}, m_s) d\vec{x} = \int \psi^*(\boldsymbol{\xi}) \psi'(\boldsymbol{\xi}) d\boldsymbol{\xi}$$

Normalization: $\int |\psi_{\uparrow}(\vec{x})|^2 d\vec{x} + \int |\psi_{\downarrow}(\vec{x})|^2 d\vec{x} = 1$

Probability density for particle at given position and spin:

$$\mathbf{p}_{\psi}(\vec{x} \uparrow) = |\mathbf{a}_{\psi}(\vec{x} \uparrow)|^2 = |\langle \vec{x} \uparrow | \psi \rangle|^2 = \left| \int (\delta(\vec{x}' - \vec{x}), 0) \begin{pmatrix} \psi_{\uparrow}(\vec{x}') \\ \psi_{\downarrow}(\vec{x}') \end{pmatrix} d\vec{x}' \right|^2 = |\psi_{\uparrow}(\vec{x})|^2 = |\psi(\vec{x}, +\frac{1}{2})|^2$$

$$\mathbf{p}_{\psi}(\vec{x} \downarrow) = |\mathbf{a}_{\psi}(\vec{x} \downarrow)|^2 = |\langle \vec{x} \downarrow | \psi \rangle|^2 = \left| \int (0, \delta(\vec{x}' - \vec{x})) \begin{pmatrix} \psi_{\uparrow}(\vec{x}') \\ \psi_{\downarrow}(\vec{x}') \end{pmatrix} d\vec{x}' \right|^2 = |\psi_{\downarrow}(\vec{x})|^2 = |\psi(\vec{x}, -\frac{1}{2})|^2$$

► **Coordinate-spin entanglement:** Almost all spinor states are entangled. A factorized state has a special structure $\boldsymbol{\psi}(\vec{x}) = \psi(\vec{x})(\alpha_1 |\uparrow\rangle + \alpha_2 |\downarrow\rangle) = \psi(\vec{x}) \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}$

► **Application in a which-path version of the double-slit experiment**

Consider a modification of the double-slit experiment (see Intro.) such that


both slits are equipped with spin polarizers: $\begin{cases} \text{A polarizes electron in direction } \uparrow \\ \text{B polarizes electron in direction } \downarrow \end{cases}$

⇒ paths through slits A & B are distinguishable

The electron's state inside the interferometer is $\boldsymbol{\psi}(\vec{x}) = \alpha \psi_A(\vec{x}) |\uparrow\rangle + \beta \psi_B(\vec{x}) |\downarrow\rangle$, where ψ_A, ψ_B are spatial wavefunctions for particles propagating from the respective slit and α, β some coefficients.

Probability density to detect the electron at place

\vec{x} reads $\mathbf{p}_{\psi}(\vec{x}) = \langle \psi | \hat{P}_{\vec{x}} | \psi \rangle$, where $\hat{P}_{\vec{x}} = |\vec{x} \uparrow\rangle \langle \vec{x} \uparrow| + |\vec{x} \downarrow\rangle \langle \vec{x} \downarrow|$ is the projector to the subspace of \mathcal{H} (more precisely $\overline{\mathcal{H}}$) spanned by vectors $|\vec{x}\rangle |\uparrow\rangle$ and $|\vec{x}\rangle |\downarrow\rangle$:



$$\begin{aligned}
\mathbf{p}_\psi(\vec{x}) &= \underbrace{\psi_A^*(\vec{x})}_{\psi_A^*(\vec{x})} \underbrace{1}_{0} + \beta^* \underbrace{\psi_B^*(\vec{x})}_{\psi_B^*(\vec{x})} \underbrace{0}_{1} \left(\alpha \underbrace{\langle \vec{x} | \psi_A \rangle}_{\psi_A(\vec{x})} \underbrace{\langle \uparrow | \uparrow \rangle}_{0} + \beta \underbrace{\langle \vec{x} | \psi_B \rangle}_{\psi_B(\vec{x})} \underbrace{\langle \uparrow | \downarrow \rangle}_{1} \right) + \\
&\quad \left(\alpha^* \underbrace{\langle \psi_A | \vec{x} \rangle}_{\psi_A(\vec{x})} \underbrace{\langle \uparrow | \downarrow \rangle}_{0} + \beta^* \underbrace{\langle \psi_B | \vec{x} \rangle}_{\psi_B(\vec{x})} \underbrace{\langle \downarrow | \downarrow \rangle}_{1} \right) \left(\alpha \underbrace{\langle \vec{x} | \psi_A \rangle}_{\psi_A(\vec{x})} \underbrace{\langle \downarrow | \uparrow \rangle}_{0} + \beta \underbrace{\langle \vec{x} | \psi_B \rangle}_{\psi_B(\vec{x})} \underbrace{\langle \downarrow | \downarrow \rangle}_{1} \right) + \\
&= |\alpha\psi_A(\vec{x})|^2 + |\beta\psi_B(\vec{x})|^2 \Rightarrow \text{no interference appears (the same holds if paths} \\
&\quad \text{A \& B are recorded by two } \textit{perpendicular} \text{ states of } \textit{any} \text{ spectator system)}
\end{aligned}$$

◀ Historical remark

1922: O. Stern & W. Gerlach observe the first indication of spin

1924: Wolfgang Pauli introduces “two-valued quantum degree of freedom” and formulates the exclusion principle (see below), in 1927 he introduces spinors

1925: R. Kronig and G. Uhlenbeck & S. Goudsmit provide an interpretation of spin in terms of intrinsic rotation (refused at that time)

■ Two or more distinguishable structureless particles with spin $\frac{1}{2}$

We are ready to construct state spaces for collections of particles. At first we assume that the particles are of different types—*distinguishable*. We assume N particles with spin $\frac{1}{2}$, but the same procedure can be applied regardless of spin.

► $\mathcal{H}_1, \mathcal{H}_2, \dots, \mathcal{H}_N =$ Hilbert spaces of individual particles: $\mathcal{H}_i = \mathcal{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^2$

$\mathcal{H}^{(N)} \equiv \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$	Wavefunction of a general state $ \Psi\rangle \in \mathcal{H}^{(N)}$
	$\Psi(\underbrace{\vec{x}_1, m_1}_{\xi_1}, \underbrace{\vec{x}_2, m_2}_{\xi_2}, \dots, \underbrace{\vec{x}_N, m_N}_{\xi_N}) = \underbrace{\langle \xi_1 \dots \xi_N \Psi \rangle}_{\alpha_\Psi(\xi_1 \dots \xi_N)}$
Scalar product: $\langle \Psi \Psi' \rangle \equiv$	
$\sum_{m_1} \dots \sum_{m_N} \int \dots \int \Psi^*(\vec{x}_1, m_1 \dots \vec{x}_N, m_N) \Psi'(\vec{x}_1, m_1 \dots \vec{x}_N, m_N) d\vec{x}_1 \dots d\vec{x}_N$	N -particle
$= \int \dots \int \Psi^*(\xi_1 \dots \xi_N) \Psi'(\xi_1 \dots \xi_N) d\xi_1 \dots d\xi_N$	amplitude

► **Multidimensional entanglement:** Almost all states exhibit all kinds of entanglement (coordinate-coordinate, spin-spin, and coordinate-spin) of different particles and coordinate-spin entanglement of identical particles

► Probability expressions

Wavefunction $\Psi(\xi_1 \dots \xi_N)$ lives in the multidimensional configuration space containing generalized coordinates $\xi_i \equiv (\vec{x}_i, m_i)$ of all particles. It contains complete information on mutual correlations between particles and allows one to extract any kind of probability distribution in the generalized coordinate space:

(a) **Joint probability** density to find particles $\# (1, 2 \dots N)$ at $(\xi_1, \xi_2 \dots \xi_N)$

$$\mathbf{p}_\Psi(\xi_1 \dots \xi_N) = |\alpha_\Psi(\xi_1 \dots \xi_N)|^2 = \langle \Psi | \xi_1 \dots \xi_N \rangle \langle \xi_1 \dots \xi_N | \Psi \rangle = |\Psi(\xi_1 \dots \xi_N)|^2$$

Normalization: $\int \dots \int \mathbf{p}_\Psi(\xi_1 \dots \xi_N) d\xi_1 \dots d\xi_N = 1$

(b) **Integrated probability** $\mathbf{p}_\Psi(X)$ of a property defined by $(\xi_1 \dots \xi_N) \in X$, where X is a certain domain in the multidimensional configuration space:

$$\mathbf{p}_\Psi(X) = \int \dots \int \chi_X(\xi_1 \dots \xi_N) \mathbf{p}_\Psi(\xi_1 \dots \xi_N) d\xi_1 \dots d\xi_N \quad \text{where } \chi_X = \begin{cases} 1 & \text{for } (\xi_1 \dots \xi_N) \in X \\ 0 & \text{for } (\xi_1 \dots \xi_N) \notin X \end{cases}$$

Example: property X identified with the subset of the config. space in which particle #1 is localized in a state between $|\xi\rangle$ and $|\xi+d\xi\rangle$. Probability $\mathbf{p}_\Psi(X) = \mathbf{p}_\Psi(\xi_1 = \xi)d\xi$, where the density: $\mathbf{p}_\Psi(\xi_1 = \xi) = \underbrace{\int \dots \int}_{N-1} \mathbf{p}_\Psi(\xi, \xi_2, \dots, \xi_N) d\xi_2 \dots d\xi_N$

(c) **Single-particle probability density** to find *any* of the N particles at ξ

$$\mathbf{p}_\Psi(\xi) = \frac{1}{N} \sum_{i=1}^N \underbrace{\int \dots \int}_{N-1} \mathbf{p}_\Psi(\xi_1 \dots \xi_{i-1} \underbrace{\xi}_i \xi_{i+1} \dots \xi_N) d\xi_1 \dots d\xi_{i-1} d\xi_{i+1} \dots d\xi_N$$

Normalization $\int \mathbf{p}_\Psi(\xi) d\xi = 1$. This density is determined from the expression $N\mathbf{p}_\Psi(\xi)d\xi = \sum_{n=0}^N n\mathbf{p}_\Psi(n, \xi, d\xi) \equiv \varrho_\Psi(\xi)d\xi$, where $\mathbf{p}_\Psi(n, \xi, d\xi)$ is the integrated probability for finding any $n \leq N$ particles in states between $|\xi\rangle$ and $|\xi+d\xi\rangle$. So $\varrho_\Psi(\xi) = N\mathbf{p}_\Psi(\xi)$ normalized as $\int \varrho_\Psi(\xi) d\xi = N$ is an **average particle concentration** (not accounting for particle identity) at generalized coordinate ξ .

■ Two indistinguishable particles

In quantum physics, if some particles are really the same, we cannot define any kind of their individuality. In particular, we cannot assign to these particles any intrinsic names/numbers as there exists no property that would enable us to recognize whether a given particle is “Fred” or “Bruno”. Consider two electrons localized at distinct places A and B . We can name them “the electron at place A ” and “the electron at place B ”; these are two distinguished single-particle states. However, it is impossible to say whether the present electron at place A is the same electron as the one observed at place A some time ago.

► The Hilbert space of two indistinguishable particles is constructed from the Hilbert space $\mathcal{H}^{(2)} \equiv \mathcal{H}_1 \otimes \mathcal{H}_2 \ni |\Psi\rangle \equiv \Psi(\xi_1, \xi_2)$ of two formally distinguishable (though physically identical) particles #1 & #2. We introduce in this space the

particle exchange operator: $\hat{E}_{1\leftrightarrow 2}\Psi(\xi_1, \xi_2) = \Psi(\xi_2, \xi_1) \quad \hat{E}_{1\leftrightarrow 2}^2 = \hat{I}$

$\hat{E}_{1\leftrightarrow 2}$ exchanges states of particles #1 & #2 in the expansion of $|\Psi\rangle$ in any factorized basis: $|\Psi\rangle = \sum_{ij} \alpha_{ij} |\phi_i\rangle_1 |\phi_j\rangle_2 \Rightarrow \hat{E}_{1\leftrightarrow 2} |\Psi\rangle = \sum_{ij} \alpha_{ij} |\phi_j\rangle_1 |\phi_i\rangle_2$

► For indistinguishable particles we require that the exchange affects only the overall phase of the state since it is physically irrelevant, so $\hat{E}_{1\leftrightarrow 2} |\Psi\rangle = e^{i\varphi} |\Psi\rangle$ with $\varphi \in \mathbb{R}$, and that two subsequent exchanges yield the original state: $e^{2i\varphi} = 1$

$$\Rightarrow \begin{cases} \varphi = 0 & \Psi(\xi_1, \xi_2) = +\Psi(\xi_2, \xi_1) & \text{symmetric} & \text{for bosons} \\ \varphi = \pi & \Psi(\xi_1, \xi_2) = -\Psi(\xi_2, \xi_1) & \text{antisymmetric} & \text{for fermions} \end{cases}$$

The two possibilities of phase φ define two fundamental types of elementary particles in nature: bosons (with exchange-symmetric wave functions) and fermions (with exchange-antisymmetric wave functions). It turns out (proof given only

in the relativistic QM) that particles with spin $\frac{1}{2}$ (or spins $\frac{3}{2}, \frac{5}{2}, \dots$) are fermions, while those with no spin (spin 0 or spins 1, 2, ...) are bosons.

► Any two-body wavefunction can be uniquely decomposed into the symmetric & antisymmetric parts that belong to distinct subspaces of $\mathcal{H}^{(2)}$:


$$\Psi(\xi_1, \xi_2) = \underbrace{\frac{1}{2}[\Psi(\xi_1, \xi_2) + \Psi(\xi_2, \xi_1)]}_{\hat{P}_+ \Psi(\xi_1, \xi_2)} + \underbrace{\frac{1}{2}[\Psi(\xi_1, \xi_2) - \Psi(\xi_2, \xi_1)]}_{\hat{P}_- \Psi(\xi_1, \xi_2)}$$

\hat{P}_+ and \hat{P}_- are projectors to the symmetric and antisymmetric subspaces

$$\hat{P}_+ + \hat{P}_- = \hat{I} \quad \Rightarrow \quad \boxed{\mathcal{H}^{(2)} = \mathcal{H}_+^{(2)} \oplus \mathcal{H}_-^{(2)}}$$

General form of decomposition for $|\Psi\rangle = \sum_{ij} \alpha_{ij} |\phi_i\rangle_1 |\phi_j\rangle_2$:

$$\hat{P}_\pm |\Psi\rangle = \sum_{ij} \alpha_{ij} \frac{1}{2} [|\phi_i\rangle_1 |\phi_j\rangle_2 \pm |\phi_j\rangle_1 |\phi_i\rangle_2]$$



► **Pauli principle:** $\hat{P}_- |\psi\rangle_1 |\psi\rangle_2 = 0$ (states $|\psi\rangle_1 |\psi\rangle_2$ are \perp to subspace $\mathcal{H}_-^{(2)}$)

\Rightarrow Two (or more) fermions cannot occur in the same single-particle state. Each single-particle state can be occupied at most by one fermion. This has tremendous consequences for the structure of matter! Without Pauli principle, the world would be a boring place (probably with no bored creature present).

► Interference effects caused by indistinguishability

Two **distinguishable** particles in a factorized state: $\Psi(\xi_1, \xi_2) = \psi_1(\xi_1)\psi_2(\xi_2)$

Joint probability density: $\mathbf{p}_\Psi(\xi_1, \xi_2) = \mathbf{p}_1(\xi_1)\mathbf{p}_2(\xi_2)$

Single-particle prob. density: $\mathbf{p}_\Psi(\xi) = \frac{1}{2}[\mathbf{p}_1(\xi) + \mathbf{p}_2(\xi)] \quad \Rightarrow$ no interference

Here $\mathbf{p}_i(\xi_i) \equiv \mathbf{p}_{\psi_i}(\xi_i) = |\psi_i(\xi_i)|^2$ is prob. density associated with particle $\#i = 1, 2$

Two **indistinguishable** particles: $\hat{P}_\pm \Psi(\xi_1, \xi_2) \propto [\psi_1(\xi_1)\psi_2(\xi_2) \pm \psi_1(\xi_2)\psi_2(\xi_1)]$


Joint: $\mathbf{p}_\Psi(\xi_1, \xi_2) \propto \mathbf{p}_1(\xi_1)\mathbf{p}_2(\xi_2) + \mathbf{p}_1(\xi_2)\mathbf{p}_2(\xi_1) \pm 2\text{Re}[\psi_1(\xi_1)\psi_2^*(\xi_1)\psi_1^*(\xi_2)\psi_2(\xi_2)]$

Single-particle: $\mathbf{p}_\Psi(\xi) \propto \mathbf{p}_1(\xi) + \mathbf{p}_2(\xi) \pm 2\text{Re}[\langle \psi_1 | \psi_2 \rangle \psi_1^*(\xi)\psi_2(\xi)]$

The state $\hat{P}_\pm \Psi(\xi_1, \xi_2)$ is entangled and this immediate consequence of particle indistinguishability creates interference effects in both probability densities $\mathbf{p}_\Psi(\xi_1, \xi_2)$ & $\mathbf{p}_\Psi(\xi)$.

However, the interference is significant only if the states $\psi_1(\xi)$ & $\psi_2(\xi)$ have a sufficient overlap. No interference effects are observed

e.g. for very distant particles or for particles with opposite spins (\Rightarrow entanglement of electrons in different galaxies, for instance, is practically unmeasurable).



■ Many indistinguishable particles

It is straightforward (but more laborious) to generalize the above results to $N > 2$ indistinguishable particles. In short, particle permutations are decomposed into

pairwise exchanges and the states of identical bosons (fermions) are identified with symmetric (antisymmetric) subspaces with respect to these exchanges. A general theory of bosonic & fermionic systems will be elaborated in Sec. 14.

► N distinguishable particles: $\Psi(\xi_1, \dots, \xi_N) \equiv |\Psi\rangle \in \mathcal{H}^{(N)} \equiv \otimes_{k=1}^N \mathcal{H}_k$

Factorized basis: $|\phi_{i_1}\rangle_1 |\phi_{i_2}\rangle_2 \dots |\phi_{i_N}\rangle_N \equiv |\Phi_{i_1 i_2 \dots i_N}\rangle$ with $i_k = 1, 2, 3, \dots$

► Particle exchanges and permutations

Exchange operators


$$\hat{E}_{k \rightleftharpoons l} |\Phi_{i_1 \dots i_k \dots i_l \dots i_N}\rangle = |\Phi_{i_1 \dots i_l \dots i_k \dots i_N}\rangle$$

Permutation operators

$$\hat{\mathcal{E}}_\pi |\Phi_{i_1 i_2 \dots i_N}\rangle = |\Phi_{i_{k_1^\pi} i_{k_2^\pi} \dots i_{k_N^\pi}}\rangle$$

Permutation $(k_1^\pi, k_2^\pi, \dots, k_N^\pi)$ is an arbitrary reordering of the original sequence $(1, 2, \dots, N)$. Index $\pi = 1, \dots, N!$ is the permutation identifier and number $k_i^\pi \in \{1, 2, \dots, N\}$ stands for the i th term of the π th reordered sequence. For example $(1, 2, 3) \rightarrow (1, 2, 3), (3, 1, 2), (2, 3, 1), (1, 3, 2), (3, 2, 1), (2, 1, 3)$ for $N=3$. Any permutation $\hat{\mathcal{E}}_\pi$ can be written as a product of exchanges $\hat{E}_{k \rightleftharpoons l}$. The factorization of a given $\hat{\mathcal{E}}_\pi$ is not unique, but all factorizations have either even or odd number of exchanges. This defines even & odd permutations.

Permutation sign $\sigma_\pi = \begin{cases} + & \text{for even permutation} \\ - & \text{for odd permutation} \end{cases}$



► Hilbert space decomposition

$$\mathcal{H}^{(N)} = \mathcal{H}_+^{(N)} \oplus \dots \oplus \mathcal{H}_-^{(N)}$$

The dots represent subspaces with mixed symmetries, while $\mathcal{H}_+^{(N)}$ and $\mathcal{H}_-^{(N)}$ are fully symmetric and antisymmetric subspaces satisfying:

$$\hat{\mathcal{E}}_\pi |\Psi\rangle = +|\Psi\rangle \quad \forall |\Psi\rangle \in \mathcal{H}_+^{(N)}, \quad \hat{\mathcal{E}}_\pi |\Psi\rangle = \sigma_\pi |\Psi\rangle \quad \forall |\Psi\rangle \in \mathcal{H}_-^{(N)}$$

$$\hat{P}_+^{(N)} = \frac{1}{N!} \sum_{\pi=1}^{N!} \hat{\mathcal{E}}_\pi$$

$$\hat{P}_-^{(N)} = \frac{1}{N!} \sum_{\pi=1}^{N!} \sigma_\pi \hat{\mathcal{E}}_\pi$$

projectors to $\mathcal{H}_+^{(N)}$ & $\mathcal{H}_-^{(N)}$

A sketch of proof: The fact that $\hat{P}_\pm |\Psi\rangle \in \mathcal{H}_\pm^{(N)}$ follows from the closure relations of permutations: (i) for any two permutations π, π' we have $\hat{\mathcal{E}}_\pi \hat{\mathcal{E}}_{\pi'} = \hat{\mathcal{E}}_{\pi''}$, where π'' denotes another permutation satisfying $\sigma_{\pi''} = \sigma_\pi \sigma_{\pi'}$, (ii) if $\{\hat{\mathcal{E}}_{\pi'}\}_{\pi'=1}^{N!}$ represents a complete set of permutations, so does $\{\hat{\mathcal{E}}_\pi \hat{\mathcal{E}}_{\pi'}\}_{\pi'=1}^{N!}$ for any fixed $\hat{\mathcal{E}}_\pi$. In this way we can show that $\hat{\mathcal{E}}_\pi \hat{P}_+^{(N)} |\Psi\rangle = \hat{P}_+^{(N)} |\Psi\rangle$ and $\hat{\mathcal{E}}_\pi \hat{P}_-^{(N)} |\Psi\rangle = \sigma_\pi \hat{P}_-^{(N)} |\Psi\rangle$. Relations (i) and (ii) also imply that $\sum_\pi \sum_{\pi'} \hat{\mathcal{E}}_\pi \hat{\mathcal{E}}_{\pi'} = N! \sum_{\pi''} \hat{\mathcal{E}}_{\pi''}$ and $\sum_\pi \sum_{\pi'} \sigma_\pi \hat{\mathcal{E}}_\pi \sigma_{\pi'} \hat{\mathcal{E}}_{\pi'} = N! \sum_{\pi''} \sigma_{\pi''} \hat{\mathcal{E}}_{\pi''}$, which prove that $(\hat{P}_\pm^{(N)})^2 = \hat{P}_\pm^{(N)}$.

Hilbert space for N identical particles is $\begin{cases} \mathcal{H}_+^{(N)} & \text{for bosons} \\ \mathcal{H}_-^{(N)} & \text{for fermions} \end{cases}$

$\hat{P}_+^{(N)} + \hat{P}_-^{(N)} \neq \hat{I}$ for $N > 2$: the rest of the space contains mixed symmetry subspaces (corresponding e.g. to mixtures of several types of identical particles)

► Expression of a basis in the fermionic space:

$$\hat{P}_-^{(N)} \underbrace{\left[|\phi_1\rangle_1 |\phi_2\rangle_2 \dots |\phi_N\rangle_N \right]}_{|\Phi_{12\dots N}\rangle} = \frac{1}{N!} \text{Det} \begin{pmatrix} |\phi_1\rangle_1 & |\phi_1\rangle_2 & \dots & |\phi_1\rangle_N \\ |\phi_2\rangle_1 & |\phi_2\rangle_2 & \dots & |\phi_2\rangle_N \\ \vdots & & & \vdots \\ |\phi_N\rangle_1 & |\phi_N\rangle_2 & \dots & |\phi_N\rangle_N \end{pmatrix} \quad \text{Slater determinant}$$

Normalization: The above projected state is not normalized. For the normalization coefficient we calculate $\langle \hat{P}_-^{(N)} \Phi_{12\dots N} | \hat{P}_-^{(N)} \Phi_{12\dots N} \rangle = \langle \Phi_{12\dots N} | (\hat{P}_-^{(N)})^2 \Phi_{12\dots N} \rangle = \langle \Phi_{12\dots N} | \hat{P}_-^{(N)} \Phi_{12\dots N} \rangle = 1/N!$ (the 2nd eq. follows from the hermiticity of projectors, see Sec. 2a, the last from the fact that any nontrivial permutation of $|\phi_1\rangle_1 \dots |\phi_N\rangle_N$, where all $|\phi_i\rangle$ s are mutually different, yields zero overlap with the original state) \Rightarrow the normalized state reads $\sqrt{N!} \hat{P}_-^{(N)} |\Phi_{12\dots N}\rangle = \frac{1}{\sqrt{N!}} \text{Det}(\dots)$

Notes: (a) Analogous expression for bosons can be formally written with “Det” replaced by a symbol denoting the exchange-symmetric sum of permutations. (b) Slater-determinant or analogous symmetrized states originate from *factorized* states in the space of distinguishable particles. They carry just a minimal unavoidable entanglement caused by indistinguishability of particles. These states form a basis in $\mathcal{H}_-^{(N)}$ or $\mathcal{H}_+^{(N)}$, so a general N -body fermionic or bosonic state can be expressed as a *superposition* of these basis states.

◀ Historical remark

1924: S.N. Bose derives Planck blackbody law from indistinguishability of photons
 1924: Wolfgang Pauli formulates the exclusion principle to explain periodic table
 1926: Werner Heisenberg and Paul Dirac relate Pauli principle to antisymmetric wavefunctions and Bose-Einstein statistics to symmetric wavefunctions. Dirac and Enrico Fermi derive statistical law for “fermions”
 1927: D. Hartree & V. Fock derive approximation for atomic N -electron wavefunctions, in 1929 J. Slater facilitates the description by using the determinant
 1939-50: M. Fierz, W. Pauli, J. Schwinger provide proofs (within the relativistic quantum theory) of the general theorem relating the “type of statistics” to spin

■ Systems with unbounded number of particles

We come to many-particle systems in which the particle number is not fixed. One can think of an exchange of particles with a bath. More fundamentally, if the special relativity is applied to processes involving elementary particles, the number of particles (the sum of their rest masses) is not conserved. Particles can be repeatedly created and annihilated, conserving only the total energy \Leftrightarrow mass of the system. It turns out that considering no upper bound on the

particle number we leave the safe harbor of separable Hilbert spaces and face the limitless ocean of continuum. This is a transition to the field theory. Work with the Fock space within the nonrelativistic QM will be practiced in Secs. 14 & 15.


► Fock space

Sum of spaces for all particle numbers $N = 0, 1, 2, 3, \dots$

$$\mathcal{H} \equiv \underbrace{\mathcal{H}^{(0)}}_{\text{vacuum state } |0\rangle} \oplus \underbrace{\mathcal{H}^{(1)}}_{1 \text{ particle}} \oplus \underbrace{\mathcal{H}^{(2)}}_{2 \text{ particles}} \oplus \dots \oplus \underbrace{\mathcal{H}^{(N)}}_{N \text{ particles}} \oplus \dots$$

This applies for distinguishable/indistinguishable particles of the same type:

$$\mathcal{H}_{\bullet}^{(N)} \equiv \begin{cases} \mathcal{H}_{+}^{(N)} & \text{indistinguishable bosons} \\ \mathcal{H}_{-}^{(N)} & \text{indistinguishable fermions} \\ \mathcal{H}^{(N)} & \text{distinguishable particles} \end{cases}$$



► Separability versus non-separability

In nonrelativistic QM it is assumed that the actual number of particles N is *unlimited* but *finite*. The set of basis vectors subject to this constraint is countable and such Fock space is **separable**.

However, the *closure* of the Fock space including $\mathcal{H}_{\bullet}^{(\infty)}$ is **non-separable**. Reasoning: basis states $|\Phi_{i_1 i_2 \dots}\rangle \equiv |\phi_{i_1}\rangle_1 |\phi_{i_2}\rangle_2 \dots$ for $N=\infty$ are specified by an infinite number of integer indices i_1, i_2, \dots identifying basis states of individual particles. This set is uncountable in analogy to real numbers (infinite sequences of digits; see Cantor's "diagonal slash" argument).

◀ Historical remark

1932: Vladimir Fock introduced the space for indefinite particle number

1958: Paul Dirac relates the Fock space to field quantization & continuum problems

■ Artificial systems (qubits...)

Since recently, various artificial quantum systems are assembled in the laboratory to be harnessed in potential applications of quantum information technologies (Sec. 9). Such systems (formed by ensembles of trapped atoms, nuclear spins, superconducting circuits etc.) are designed so that they allow for controlled manipulations and show minimal sensitivity to external perturbations. The specific physical content of individual states is not essential (this being just an "engineering" issue) and the only focus is set to the mathematical properties of complex superpositions of arbitrary basis states in the finite Hilbert space.

► **Qubit:** any system with the 2-dimensional Hilbert space $\boxed{\mathcal{H} = \mathbb{C}^2}$ can be considered as a quantum analog of classical bit. The basis of \mathcal{H} (formed by two selected states of the underlying system) is denoted as $\{|0\rangle, |1\rangle\}$.

General normalized states of the qubit are mapped to the

$$|\psi\rangle = \alpha_0|0\rangle + \alpha_1|1\rangle = \overbrace{e^{i\varphi_0}}^{\text{irrel. global phase}} \left(\cos \frac{\vartheta}{2} |0\rangle + e^{i(\varphi_1 - \varphi_0)} \sin \frac{\vartheta}{2} |1\rangle \right)$$

unit sphere (points with spherical angles ϑ, φ), where classical logical states correspond to the north ($|0\rangle$) and south ($|1\rangle$) poles.

► **Qudit**: a generalization to any higher dimension d (e.g., qutrit for $d = 3$ etc.). Hilbert space $\mathcal{H} = \mathbb{C}^d \equiv \text{Span}\{|0\rangle, |1\rangle, \dots, |d-1\rangle\}$. A general normalized state $|\psi\rangle = \sum_{k=0}^{d-1} \alpha_k |k\rangle$ is determined (up to the global phase) by $2(d-1)$ real parameters (e.g., by $d-1$ hyperspherical angles and $d-1$ relative phase angles).

► **System of N qubits** (Hilbert space of a quantum computer)

$$\mathcal{H} = \bigotimes_{i=1}^N \mathcal{H}_i \equiv \text{Span}\left\{ \underbrace{|l_1\rangle_1 |l_2\rangle_2 \dots |l_N\rangle_N}_{|x\rangle} \right\}_{(l_1, l_2, \dots, l_N) = (0, 0, \dots, 0)}^{(1, 1, \dots, 1)} \equiv \text{Span}\{|x\rangle\}_{x=0}^{2^N-1} \quad d = 2^N$$

Factorized basis vectors built from states $|l_i\rangle_i \equiv |0\rangle_i$ or $|1\rangle_i$ can be enumerated by $x = 0, \dots, 2^N - 1$ so that $l_1 l_2 \dots l_N$ is the binary representation of x .

General state of the system describes a **quantum register** that can carry integers $x \in \{0, 1, \dots, 2^N - 1\}$ as well as all their superpositions with any coefficients $\alpha_x \in \mathbb{C}$.

$$|\Psi\rangle = \sum_{x=0}^{2^N-1} \alpha_x |x\rangle$$

Although the initial state $|\Psi_0\rangle$ of various computational procedures is factorized, $|\Psi_0\rangle = \frac{1}{\sqrt{2^N}} \sum_{x=0}^{2^N-1} |x\rangle = \prod_{i=1}^N \frac{1}{\sqrt{2}} (|0\rangle_i + |1\rangle_i)$ the space \mathcal{H} supports all kinds of *bi- & multi-partite entanglement*. Splitting the whole register to 2 subregisters A and B with n and $N-n$ qubits, respectively, $\mathcal{H}_A = \bigotimes_{i=1}^n \mathcal{H}_i = \text{Span}\{|x\rangle_A\}_{x=0}^{2^n-1}$, $\mathcal{H}_B = \bigotimes_{i=n+1}^N \mathcal{H}_i = \text{Span}\{|x'\rangle_B\}_{x'=0}^{2^{N-n}-1}$, so $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, we can express a general state $\in \mathcal{H}$ in the form

$$|\Psi\rangle = \sum_{x=0}^{2^n-1} \sum_{x'=0}^{2^{N-n}-1} \alpha_{xx'} |x\rangle_A |x'\rangle_B \quad \alpha_{xx'} \in \mathbb{C}$$

It is **almost always entangled!**

The space \mathcal{H} can be decomposed to subspaces with various **exchange symmetries** of individual qubit states. Consider concrete splitting s to 2 subregisters with n and $N-n$ qubits (e.g., $s = \{1, 3, 4\}\{2, 5\}$ for $N = 5$) and define the subspace $\mathcal{H}_s^{(N, n)}$ such that $\hat{E}_{i=j} |\Psi\rangle = \pm |\Psi\rangle$ for any $|\Psi\rangle \in \mathcal{H}_s^{(N, n)}$, with $\begin{cases} + & \text{for } i, j \text{ in the same subregister,} \\ - & \text{for } i, j \text{ in different subregisters.} \end{cases}$ Let $r(N, n)$ is the number such (nonequivalent) subspaces. It can be shown that $\mathcal{H} = \bigoplus_{n=\lceil N/2 \rceil}^N \bigoplus_{s=1}^{r(N, n)} \mathcal{H}_s^{(N, n)}$

◀ Historical remark

1980's: P. Benioff, R. Feynman, D. Deutsch and others pioneer the idea of using controllable quantum systems for simulation and computation purposes

1995-present: construction of increasingly complex controllable quantum devices

Requirement 2: Hermiticity

$$\langle \psi_1 | \hat{A} \psi_2 \rangle = \langle \hat{A} \psi_1 | \psi_2 \rangle = \langle \psi_2 | \hat{A} \psi_1 \rangle^*$$

$$\langle \psi | \hat{A}^n \psi \rangle \in \mathbb{R}$$

Hermiticity is sufficient (though not necessary) condition for the statistical moments being **real numbers** (as needed)

\Rightarrow the matrices associated with \hat{A} satisfy: $A_{ij} = A_{ji}^*$ for $i \neq j$ and $A_{ii} \in \mathbb{R}$

► Some mathematical definitions

Definition domain: Operator \hat{A} is defined for $|\psi\rangle \in \text{Def}(\hat{A}) \subseteq \mathcal{H}$

For physics purposes it often suffices if

$$\text{Def}(\hat{A}) \equiv \text{a dense subset } \underline{\mathcal{H}} \subset \mathcal{H}$$

Operator norm: $\|\hat{A}\|^2 \equiv \text{Sup} \left\{ \frac{\langle \hat{A} \psi | \hat{A} \psi \rangle}{\langle \psi | \psi \rangle} \right\}_{|\psi\rangle \in \text{Def}(\hat{A})}$ (cf. rigged Hilbert space, Sec. 1a)

$\|\hat{A}\| < \infty$ for **bounded** operators, $\|\hat{A}\| = \infty$ for **unbounded** operators

Hermitian adjoint operator \hat{A}^\dagger is an operator satisfying the condition:

$$\langle \psi_1 | \hat{A} \psi_2 \rangle = \langle \hat{A}^\dagger \psi_1 | \psi_2 \rangle = \langle \psi_2 | \hat{A}^\dagger \psi_1 \rangle^* \quad \begin{cases} \forall |\psi_2\rangle \in \text{Def}(\hat{A}) \\ \forall |\psi_1\rangle \in \text{Def}(\hat{A}^\dagger) \supseteq \text{Def}(\hat{A}) \end{cases}$$

$$\hat{A}^\dagger = \begin{pmatrix} A_{11}^* & A_{21}^* & \dots \\ A_{12}^* & A_{22}^* & \\ \vdots & & \ddots \end{pmatrix} \equiv \hat{A}^{\text{T}*}$$

matrix representation of Hermitian adjoint operator
(= transpose & complex conjugate matrix)

Adjoint of a product: $(\hat{A}_1 \hat{A}_2)^\dagger = \hat{A}_2^\dagger \hat{A}_1^\dagger$ and similarly for multiple products

$$\langle \psi_1 | \hat{A}_1 \hat{A}_2 \psi_2 \rangle = \langle \hat{A}_1^\dagger \psi_1 | \hat{A}_2 \psi_2 \rangle = \langle \hat{A}_2^\dagger \hat{A}_1^\dagger \psi_1 | \psi_2 \rangle$$

Symmetric, selfadjoint vs. Hermitian operators

All these operators satisfy the following condition:

$$\hat{A}|\psi\rangle = \hat{A}^\dagger|\psi\rangle$$

but for different domains of vectors $|\psi\rangle$

Symmetric operator: $|\psi\rangle \in \text{Def}(\hat{A}) \subseteq \text{Def}(\hat{A}^\dagger) \subseteq \mathcal{H}$

Selfadjoint operator: $|\psi\rangle \in \text{Def}(\hat{A}) = \text{Def}(\hat{A}^\dagger) \subseteq \mathcal{H}$

Hermitian operator: $|\psi\rangle \in \text{Def}(\hat{A}) = \text{Def}(\hat{A}^\dagger) = \mathcal{H}$



These definitions are equivalent in finite-dimensional spaces

but not in ∞ -dim. spaces. Nevertheless, most textbooks including this one make use of the term “Hermitian operator” regardless of the definition domain.

► Function of operator

Physical observables are often defined as functions of other observables.

We first define a function of operator for functions of the form $f(x) = \sum_k f_k x^k$, i.e. expressible as Taylor series:

$$f(\hat{A}) \equiv \sum_k f_k \hat{A}^k$$

A more general definition will be given below

$$f_k \in \mathbb{R} \Rightarrow f(\hat{A}) \text{ Hermitian}$$

► Tensor products of operators

We will need to use operators in product spaces. Here are some constructions:

Let $\left\{ \begin{array}{l} \hat{A}_1 \text{ on } \mathcal{H}_1 \\ \hat{A}_2 \text{ on } \mathcal{H}_2 \end{array} \right\}$ be operators defined by basis actions $\left\{ \begin{array}{l} \hat{A}_1 |\phi_{1i}\rangle \equiv |\phi'_{1i}\rangle \\ \hat{A}_2 |\phi_{2j}\rangle \equiv |\phi'_{2j}\rangle \end{array} \right\}$. Then:

(a) Operator $\hat{A} = \hat{A}_1 \otimes \hat{A}_2$ on $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$ is defined by: $\hat{A}|\psi\rangle = \hat{A}[\sum_{i,j} \gamma_{ij} \overbrace{|\phi_{1i}\rangle|\phi_{2j}\rangle}^{|\Phi_{ij}\rangle}] = \sum_{i,j} \gamma_{ij} \overbrace{|\phi'_{1i}\rangle|\phi'_{2j}\rangle}^{|\Phi'_{ij}\rangle}$

(b) Possible extension of \hat{A}_1, \hat{A}_2 to $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$ is obtained from:

$$\begin{array}{l} \hat{A}_1^{(\text{ext})} = \hat{A}_1 \otimes \hat{I}_2 \equiv \hat{A}_1 \\ \hat{A}_2^{(\text{ext})} = \hat{I}_1 \otimes \hat{A}_2 \equiv \hat{A}_2 \end{array} \quad \begin{array}{l} \hat{I}_1 \equiv \text{unit op. in } \mathcal{H}_1 \\ \hat{I}_2 \equiv \text{unit op. in } \mathcal{H}_2 \end{array}$$


Similarly for multiple products $\otimes_i \mathcal{H}_i$

■ Eigenvalues and eigenvectors of Hermitian operators

The key characteristic of any operator in the Hilbert space is its spectrum of eigenvalues and the set of the corresponding eigenvectors. Not only these eigensolutions constitute a subject of an involved mathematical theory, they also play the most essential role in the formulation of quantum mechanics.

► “Dispersion-free” states and possible measurement outcomes

Consider state $|\psi_a\rangle$ in which observable A yields a “sharp” value, i.e. a single possible output a with probability $\mathbf{p}_\psi(a) = 1$. Hence the average



$\langle A \rangle_{\psi_a} = a$ and variance $\langle\langle A^2 \rangle\rangle_{\psi_a} \equiv \langle A^2 \rangle_{\psi_a} - \langle A \rangle_{\psi_a}^2 = 0$

$$\underbrace{\langle \psi_a | \hat{A}^2 \psi_a \rangle}_{\langle A^2 \rangle_{\psi_a}} - \underbrace{\langle \psi_a | \hat{A} \psi_a \rangle^2}_{\langle A \rangle_{\psi_a}^2 = a^2} = \langle \psi_a | \underbrace{\hat{A}^2 - 2a\hat{A} + a^2\hat{I}}_{(\hat{A} - a\hat{I})^2} | \psi_a \rangle = \langle (\hat{A} - a\hat{I})\psi_a | (\hat{A} - a\hat{I})\psi_a \rangle = 0$$

$$\Rightarrow (\hat{A} - a\hat{I})|\psi_a\rangle = 0 \Rightarrow \boxed{\hat{A}|\psi_a\rangle = a|\psi_a\rangle} \Rightarrow \left\{ \begin{array}{l} |\psi_a\rangle \equiv \text{eigenvector} \\ a \equiv \text{eigenvalue} \end{array} \right\} \text{ of operator } \hat{A}$$

For $\hat{A} = \hat{A}^\dagger$ the eigenvalues $a = \langle a | \hat{A} | a \rangle \in \mathbb{R}$

These considerations lead to a plausible determination of the set $\mathcal{S}(\hat{A})$ of possible measurement outcomes of observable $A \leftrightarrow \hat{A}$. We assume that *each possible outcome is associated with a state in which it is measured with certainty*, hence $\forall a \in \mathcal{S}(\hat{A})$ there \exists dispersion-free state $|\psi_a\rangle$. This leads to the postulate:

$\mathcal{S}(\hat{A}) \equiv \{ \text{possible measurement outcomes of } A \} \equiv \{ \text{eigenvalues of } \hat{A} \}$

Below we will use a “stammering” notation with $|\psi_a\rangle \equiv |a\rangle$, so:

$$\boxed{\hat{A}|a\rangle = a|a\rangle}$$

► Orthogonality of eigenvectors with different eigenvalues

$$\left. \begin{array}{l} \hat{A}|a\rangle = a|a\rangle \Rightarrow \langle a' | \hat{A} | a \rangle = a \langle a' | a \rangle \\ \hat{A}|a'\rangle = a'|a'\rangle \Rightarrow \langle a | \hat{A} | a' \rangle = \underbrace{a'}_{\langle a' | \hat{A} | a \rangle^*} \underbrace{\langle a | a' \rangle}_{\langle a' | a \rangle^*} \end{array} \right\} \begin{array}{l} \text{(valid for Hermitian operators)} \\ \Rightarrow 0 = \underbrace{(a - a')}_{\neq 0} \langle a' | a \rangle \Rightarrow \boxed{\langle a' | a \rangle = 0} \end{array}$$

\Rightarrow Different dispersion-free states (i.e., eigenstates with different eigenvalues) are perfectly distinguishable

► Degeneracy

A single eigenvalue a of \hat{A} may have *more than one linearly independent eigenvectors* $\{|a; k\rangle\}_{k=1}^n$. Due to linearity of \hat{A} , any superposition of $\{|a; k\rangle\}_{k=1}^n$ is also

an eigenvector with the same eigenvalue: $\hat{A}(\sum_{k=1}^n \alpha_k |a; k\rangle) = a \sum_{k=1}^n \alpha_k |a; k\rangle$. Hence all eigenvectors with the same eigenvalue a form a **degeneracy subspace** \mathcal{H}_a whose dimension $d_{\mathcal{H}_a} \equiv d_a \leq d_{\mathcal{H}}$ is the maximal number of linearly independent eigenvectors, i.e. the maximal size n_{\max} of the set $\{|a; k\rangle\}_{k=1}^n$. One can choose in \mathcal{H}_a an **orthonormal basis** $\{|a^{(k)}\rangle\}_{k=1}^{d_a}$ satisfying $\langle a^{(k')} | a^{(k)} \rangle = \delta_{k'k}$

► Eigensolutions for finite dimension

$$\hat{A}|a\rangle = a|a\rangle \quad \Leftrightarrow \quad (\hat{A} - a\hat{I})|a\rangle = 0 \quad \Rightarrow$$

$$\underbrace{\text{Det}(\hat{A} - a\hat{I})}_{\mathcal{P}_{\hat{A}}(a)} = 0$$

For $d_{\mathcal{H}} < \infty$ the middle relation represents a finite set of linear equations with r.h.s.=0.


Its solution exists *iff* a is a root of the **characteristic polynomial**

$$\mathcal{P}_{\hat{A}}(a) = \text{Det} \begin{pmatrix} A_{11}-a & A_{12} & \dots \\ A_{21} & A_{22}-a & \dots \\ \vdots & & \ddots \end{pmatrix}$$

of order $d_{\mathcal{H}}$. The eq. $\mathcal{P}_{\hat{A}}(a) = 0$ has $n \in \{1, 2, \dots, d_{\mathcal{H}}\}$ solutions $\{a_i\}_{i=1}^n$, all $a_i \in \mathbb{R}$ (from Hermiticity of \hat{A}).

The corresponding eigenvectors $|a_i\rangle$

satisfy $(\hat{A} - a_i\hat{I})|a_i\rangle = 0$, which for an expansion $|a_i\rangle = \sum_m \alpha_{mi} |\phi_m\rangle$ in arbitrary basis $\{|\phi_m\rangle\}_{m=1}^{d_{\mathcal{H}}}$ yields a linear set of equations for coefficients $\{\alpha_{mi}\}$. It can be solved due to the nullity of its determinant.



$$\begin{pmatrix} A_{11}-a_i & A_{12} & \dots \\ A_{21} & A_{22}-a_i & \dots \\ \vdots & & \ddots \end{pmatrix} \begin{pmatrix} \alpha_{1i} \\ \alpha_{2i} \\ \vdots \end{pmatrix} = 0$$

Theorem: For any Hermitian operator \hat{A} in \mathcal{H} , there exists $d_{\mathcal{H}}$ orthonormal eigenvectors (irrespectively of the number $1 \leq n \leq d_{\mathcal{H}}$ of eigenvalues).

Sketch of proof: For any matrix \hat{A} of dim. $d_{\mathcal{H}}$, the fundamental theorem of algebra guarantees the existence of at least one eigenvalue a and the corresponding eigenvector $|a\rangle$. The eigenvector can be normalized to $\langle a|a\rangle = 1$ since $\hat{A}|a\rangle = a|a\rangle \Rightarrow \hat{A}(\alpha|a\rangle) = a(\alpha|a\rangle) \forall \alpha \in \mathbb{C}$. Consider the orthogonal complement \mathcal{H}^\perp of $|a\rangle$, which is a $(d_{\mathcal{H}} - 1)$ dimensional subspace of \mathcal{H} such that $\langle a|\psi\rangle = 0 \forall |\psi\rangle \in \mathcal{H}^\perp$. From Hermiticity of \hat{A} we get: $\langle a|\hat{A}\psi\rangle = a\langle a|\psi\rangle$, so \mathcal{H}^\perp is invariant under the action of \hat{A} . Hence the same procedure can be repeated for \mathcal{H}^\perp , finding a new normalized eigenvector $|a'\rangle \in \mathcal{H}^\perp$. The theorem is then proven by induction.

\Rightarrow If the number of eigenvalues $n < d_{\mathcal{H}}$, some of them must be degenerate.

Dimensions of the degeneracy subspaces satisfy: $\sum_{i=1}^n d_{a_i} = d_{\mathcal{H}}$

► Completeness for finite dimension

Given any Hermitian operator \hat{A} in \mathcal{H} of dimension $d_{\mathcal{H}} < \infty$, one can introduce an **orthonormal basis** of \mathcal{H} formed by eigenvectors of \hat{A} :

Detailed notation: $\left. \begin{array}{l} \left\{ |a_i\rangle \equiv |a_i^{(1)}\rangle \text{ eigenvector for nondeg. eigenvalue } a_i \right. \\ \left. \left\{ \left\{ |a_i^{(k)}\rangle \right\}_{k=1}^{d_{a_i}} \text{ eigenvectors for degenerate eigenvalue } a_i \right. \right. \\ \left. \left. \left(\text{selected orthonormal basis of } \mathcal{H}_{a_i} \right) \right\}_{i=1}^n \right\}$

(i runs over different eigenvalues, k over different eigenvectors in a single deg. subspace)

Simplified notation: $\{a_j\}_{j=1}^{d_{\mathcal{H}}} \longleftrightarrow \{|a_j\rangle\}_{j=1}^{d_{\mathcal{H}}} \equiv \{\text{eigenvalues, some of them maybe equal}\} \longleftrightarrow \{\text{the corresponding eigenvectors, all orthonormal}\}$

Orthonormality condition:

$$\langle a_{i'}^{(k')} | a_i^{(k)} \rangle = \delta_{i'i} \delta_{k'k} \quad \langle a_{j'} | a_j \rangle = \delta_{j'j}$$

The completeness relation reads:

$$\sum_{i=1}^n \sum_{k=1}^{d_{a_i}} |a_i^{(k)}\rangle \langle a_i^{(k)}| = \hat{I}_{\mathcal{H}} = \sum_{j=1}^{d_{\mathcal{H}}} |a_j\rangle \langle a_j|$$

projector \hat{P}_{a_i}

► Diagonal representation (diagonalization)

Similarity transformation of any Hermitian matrix \hat{A} with matrix \hat{U} build from eigenvector components: $|a_j\rangle = \sum_{m=1}^{d_{\mathcal{H}}} \alpha_{mj} |\phi_m\rangle \equiv \begin{pmatrix} \alpha_{1j} \\ \alpha_{2j} \\ \vdots \end{pmatrix}$

$$\overbrace{\begin{pmatrix} A_{11} & A_{12} & \dots \\ A_{12}^* & A_{22} & \dots \\ \vdots & \ddots & \ddots \end{pmatrix}}^{\hat{A}} = \overbrace{\begin{pmatrix} \alpha_{11} & \alpha_{12} & \dots \\ \alpha_{21} & \alpha_{22} & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}}^{\hat{U}} \overbrace{\begin{pmatrix} a_1 & 0 & \dots \\ 0 & a_2 & \dots \\ \vdots & \ddots & \ddots \end{pmatrix}}^{\hat{A}_{\text{diag}}} \overbrace{\begin{pmatrix} \alpha_{11}^* & \alpha_{21}^* & \dots \\ \alpha_{12}^* & \alpha_{22}^* & \dots \\ \vdots & \ddots & \ddots \end{pmatrix}}^{\hat{U}^\dagger} \left| \begin{array}{l} \langle a_1 | \\ \langle a_2 | \\ \vdots \end{array} \right.$$

$|a_1\rangle \quad |a_2\rangle \quad \dots$

► **Example:** general 2×2 Hermitian matrix $\hat{A} = \begin{pmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{pmatrix}$ has 1 or 2 eigenvalues, but always 2 orthonorm. eigenvectors:

Characteristic polynomial: $\mathcal{P}_{\hat{A}}(a) = (A_{11} - a)(A_{22} - a) - |A_{12}|^2$
 $= a^2 - (A_{11} + A_{22})a + (A_{11}A_{22} - |A_{12}|^2)$

Roots = eigenvalues: $a_{\pm} = \frac{A_{11} + A_{22}}{2} \pm \sqrt{\left(\frac{A_{11} - A_{22}}{2}\right)^2 + |A_{12}|^2}$
 \Rightarrow degeneracy $a_+ = a_-$ iff $A_{11} = A_{22}$ and $A_{12} = 0$

Eigenvector eq.: $\begin{pmatrix} \frac{A_{11} - A_{22}}{2} \mp \sqrt{\left(\frac{A_{11} - A_{22}}{2}\right)^2 + |A_{12}|^2} & A_{12} \\ A_{12}^* & \frac{A_{22} - A_{11}}{2} \mp \sqrt{\left(\frac{A_{11} - A_{22}}{2}\right)^2 + |A_{12}|^2} \end{pmatrix} \begin{pmatrix} \alpha_{1\pm} \\ \alpha_{2\pm} \end{pmatrix} = 0$

Rows of this matrix are dependent, so the solution is determined by any of the two eqs., e.g. the first one: $\left[\frac{A_{11} - A_{22}}{2} \mp \sqrt{\left(\frac{A_{11} - A_{22}}{2}\right)^2 + |A_{12}|^2} \right] \alpha_{1\pm} + A_{12} \alpha_{2\pm} = 0$

It can be checked that $\langle a_- | a_+ \rangle = (\alpha_{1-}^*, \alpha_{2-}^*) \begin{pmatrix} \alpha_{1+} \\ \alpha_{2+} \end{pmatrix} = 0$

Counterexample: non-Hermitian matrix $\hat{A} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has only 1 eigenvector:

Characteristic polynomial: $\mathcal{P}_{\hat{A}}(a) = (1 - a)^2 \Rightarrow$ root: $a = 1$

Eigenvector eq.: $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} = 0 \Rightarrow$ single normalized eigenvector $\begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$

► **Quantization:** Already at this stage we can conclude that discrete character of the observable values of some quantities A is a consequence of the assumption that the observable values coincide with the eigenvalues a_1, a_2, \dots of the corresponding operator \hat{A} . However, the full picture cannot be drawn without considering operators in infinite-dimensional spaces.

► Eigensolutions for infinite dimension

For $d_{\mathcal{H}} = \infty$, the expression $\text{Det}(\hat{A} - a\hat{I})$ has no sense. To find solutions of the eigenvector equation $(\hat{A} - a\hat{I})|a\rangle = 0$ is much more difficult in this case. In general, an ∞ -dimensional operator \hat{A} may have both **discrete & continuous spectrum** of eigenvalues. Moreover, eigenvalues from the continuous spectrum have no eigenvectors $\in \mathcal{H}$. Note that a rigorous analysis of these issues goes beyond our present level of advancement. We will just indicate two alternative mathematical treatments: one by von Neumann, who stays within the standard Hilbert space as he allows only finite intervals of continuous eigenvalues, and one initiated by Dirac, who steps out towards the rigged Hilbert space by taking into account single eigenvalues of continuous quantities.

Example of ∞ -dim. operator with **discrete spectrum**:

$$\hat{A} = \begin{pmatrix} \boxed{0} & 0 & 0 & \dots \\ 0 & \boxed{0} & 0 & \\ 0 & 0 & \boxed{0} & \\ \vdots & & & \ddots \end{pmatrix} \Rightarrow \begin{pmatrix} \boxed{1} \\ 0 \\ 0 \\ \vdots \end{pmatrix}, \begin{pmatrix} \boxed{0} \\ \frac{1}{\sqrt{2}} \\ \pm \frac{1}{\sqrt{2}} \\ 0 \\ \vdots \end{pmatrix}, \begin{pmatrix} \boxed{0} \\ 0 \\ \frac{1}{\sqrt{2}} \\ \pm \frac{1}{\sqrt{2}} \\ \vdots \end{pmatrix}, \dots \equiv \begin{cases} \text{orthonormal} \\ \text{eigenvectors} \\ \text{corresponding} \\ \text{to eigenvalues} \\ 0, \pm 1, \pm 2, \dots \end{cases}$$

The spectrum $\mathcal{S}(\hat{A}) \equiv \mathbb{Z} \equiv \{\dots, -2, -1, 0, +1, +2, \dots\}$

The corresponding eigenvectors $|a_i\rangle \in \mathcal{H} \equiv \ell^2$

Example of ∞ -dim. operator with **continuous spectrum**:

$$\hat{A} = \begin{pmatrix} 0 & \boxed{1} & 0 & 0 & \dots \\ \boxed{1} & 0 & \boxed{1} & 0 & \\ 0 & \boxed{1} & 0 & \boxed{1} & 0 \\ 0 & 0 & \boxed{1} & 0 & \boxed{1} \\ \vdots & & & \ddots & \ddots \end{pmatrix} \Rightarrow \underbrace{\begin{pmatrix} -a & 1 & 0 & 0 & 0 & \dots \\ 1 & -a & 1 & 0 & 0 \\ 0 & 1 & -a & 1 & 0 \\ 0 & 0 & 1 & -a & 1 \\ \vdots & & \ddots & \ddots & \ddots \end{pmatrix}}_{\hat{A} - a\hat{I}} \underbrace{\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \\ \vdots \end{pmatrix}}_{|a\rangle} = 0$$

eigenvector equation:

For a finite dimension $d_{\mathcal{H}} \equiv d$ the above eigenvector equation leads to the following set of equations:

$$\left\{ \begin{array}{l} \alpha_2 = a\alpha_1 \\ \alpha_1 + \alpha_3 = a\alpha_2 \\ \alpha_2 + \alpha_4 = a\alpha_3 \\ \alpha_3 + \alpha_5 = a\alpha_4 \\ \vdots \\ \alpha_{d-2} + \alpha_d = a\alpha_{d-1} \\ \alpha_{d-1} = a\alpha_d \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{single recursive eq.} \\ \boxed{\alpha_{l-1} + \alpha_{l+1} = a\alpha_l} \\ \text{valid for } l = 1, 2, \dots, d \\ \text{with boundary} \\ \text{conditions: } \alpha_0 = \alpha_{d+1} = 0 \end{array} \right.$$

Solution for $d < \infty$: Starting from $\alpha_1 = 1$, the above set of equations yields: $\alpha_2 = a$, $\alpha_3 = (a^2 - 1)$, $\alpha_4 = (a^3 - 2a)$, ... However, the last pair of equations will not be satisfied for all values of a . It can be shown that $|\langle \alpha | \hat{A} | \alpha \rangle| \leq 2 \langle \alpha | \alpha \rangle \forall |\alpha\rangle \in \ell^2$. Hence the solution exists only for some discrete values $a_i \in [-2, +2]$ (see the figure) and the corresponding eigenvectors $|a_i\rangle$ are trivially normalizable.


Solution for $d = \infty$: In this case, the problem with the last pair of equations does not take place, so the solution exists for all $a \in (-\infty, +\infty)$. However, the

eigenvectors obtained in this way are not normalizable, i.e., $|a\rangle \notin \mathcal{H} \equiv \ell^2$. It can be shown that $|a\rangle \notin \overline{\mathcal{H}} \equiv \overline{\ell^2}$ for $|a| > 2$, so $|a\rangle$ is out of even the upper space in the Gelfand triple. On the other hand, for $|a| \leq 2$ we find $|a\rangle \in \overline{\ell^2}$.

Sketch of proof: We can solve the above recursive equation $\alpha_{l-1} + \alpha_{l+1} = a\alpha_l$ by the ansatz $\alpha_l = r^l$, which leads to the condition $r^{l-1}(1 - ar + r^2) = 0$. This yields 2 possible solutions:

$r = \frac{a}{2} \pm \sqrt{(\frac{a}{2})^2 - 1} \equiv r_{\pm}$. The boundary condition $\alpha_0 = 0$ implies $\alpha_l = (r_+^l - r_-^l)$. For $|a| \leq 2$ we have $r_{\pm} \in \mathbb{C}$. One can find $\vartheta \in [0, 2\pi)$ such that $\frac{a}{2} = \cos \vartheta$ and $r_{\pm} = \cos \vartheta \pm i \sin \vartheta = e^{\pm i\vartheta}$. Hence $\alpha_l = \sin(l\vartheta)$, which means that the components are bounded and oscillate with l and the resulting vector $|a\rangle \in \overline{\ell^2}$. In contrast, for $|a| > 2$ we have $r_{\pm} \in \mathbb{R}$ with $r_+ > 1$ and $r_- < 1$, so $|\alpha_l|$ diverges exponentially with l , which means that $|a\rangle \notin \overline{\ell^2}$.


Conclusion: Eigensolutions $\in \overline{\mathcal{H}}$ are declared to be generalized eigenvectors of an ∞ -dimensional operator \hat{A} , while those $\notin \overline{\mathcal{H}}$ are not considered as eigenvectors in any sense. Therefore, the above operator \hat{A} for $d = \infty$ has a continuous spectrum $\mathcal{S}(\hat{A}) \equiv [-2, +2]$.



► **Spectrum of a general operator in infinite dimension**

A general Hermitian operator \hat{A} can combine both discrete and continuous spectra:

$$\underbrace{\mathcal{S}(\hat{A})}_{\text{spectrum}} = \underbrace{\mathcal{D}(\hat{A})}_{\text{discrete part}} \cup \underbrace{\mathcal{C}(\hat{A})}_{\text{continuous part}}$$



Eigenvalues $a_i \in \mathcal{D}(\hat{A})$ have eigenvectors $|a_i^{(k)}\rangle \in \mathcal{H}$ where $k \in \{1, 2, \dots, d_{a_i}\}$ is a discrete degeneracy index

Eigenvalues $a \in \mathcal{C}(\hat{A})$ have eigenvectors $|a^{(k)}\rangle \in \overline{\mathcal{H}} \supset \mathcal{H}$ where $k \in \mathcal{D}_a$ is a $\left\{ \begin{array}{l} \text{discrete } (k=1, \dots, d_a) \\ \text{continuous} \end{array} \right\}$ degeneracy index

The discrete part of the spectrum fulfills the standard eigenvector relations:

Orthonormality: $\langle a_{i'}^{(k')} | a_i^{(k)} \rangle = \delta_{i'i} \delta_{k'k}$ Projectors: $\hat{P}_{a_i} = \sum_{k=1}^{d_{a_i}} |a_i^{(k)}\rangle \langle a_i^{(k)}|$

The continuous part of spectrum needs special treatment of eigenvector issues. This requires rather involved mathematics, of which we present only a rough outline.

Vážení čtenáři, právě jste dočetli ukázkou z knihy ***A Condensed Course of Quantum Mechanics*** .
Pokud se Vám ukázka líbila, na našem webu si můžete zakoupit celou knihu.